

B-ORBITS OF NILPOTENT ORDER 2 AND LINK PATTERNS

ANNA MELNIKOV

ABSTRACT. Let B_n be the group of upper-triangular invertible $n \times n$ matrices and $\mathcal{X}_n$ be the variety of strictly upper triangular $n \times n$ matrices of nilpotent order 2. B_n acts on $\mathcal{X}_n$ by conjugation. In this paper we describe geometry of orbits $\mathcal{X}_n/B_n$ in terms of link patterns.

Further we apply this description to the computations of the closures of orbital varieties of nilpotent order 2 and intersections of components of a Springer fiber of nilpotent order 2. In particular we connect our results to the combinatorics of meanders and Temperley-Lieb algebras.

1. INTRODUCTION

1.1. Let $\mathbf{M} = \mathbf{M}_n(\mathbb{C})$ be an algebra of $n \times n$ matrices over complex numbers. Let $\mathbf{G} = GL_n(\mathbb{C})$ be a general linear group. Consider its action on $\mathbf{M}$ by conjugation. For $u \in \mathbf{M}$ let $\mathcal{O}_u := \mathbf{G}.u := \{AuA^{-1} \mid A \in \mathbf{G}\}$ denote its orbit.

Let $\mathfrak{N} = \{u \in \mathbf{M} \mid u^k = 0 \text{ for some } k\}$ be the nilpotent cone of $\mathbf{M}$. Classical Jordan-Gerstenhaber theory gives a complete combinatorial description of geometry of the variety of (nilpotent) orbits $\mathcal{O}_u$ where $u \in \mathfrak{N}$ in terms of Young diagrams.

Let $\mathfrak{n} = \mathfrak{n}_n \subset \mathbf{M}$ be a subalgebra of strictly upper-triangular matrices. Let $\mathbf{B} \subset \mathbf{G}$ be a (Borel) subgroup of upper-triangular invertible matrices. Consider its action on $\mathfrak{n}$ by conjugation. For $u \in \mathfrak{n}$ let $\mathcal{B}_u := \mathbf{B}.u = \{AuA^{-1} \mid A \in \mathbf{B}\}$ denote its orbit under action of $\mathbf{B}$. In general for $n \geq 6$ the number of such orbits is infinite and there is no combinatorial theory in the spirit of Jordan-Gerstenhaber theory describing the geometry of these orbits.

However, if we take a subvariety of matrices of nilpotent order 2, that is $\mathcal{X} := \mathcal{X}_n := \{u \in \mathfrak{n} \mid u^2 = 0\}$ then the number of $\mathbf{B}$ orbits in $\mathcal{X}$ is equal to the number of involutions of the symmetric group $\mathbf{S}_n$. We show in the paper that the complete combinatorial description of geometry of $\mathbf{B}$ orbits in $\mathcal{X}$ can be obtained in terms of link patterns in the spirit of Jordan-Gerstenhaber theory.

Further we give two important application of the technique of link patterns, one in the theory of orbital varieties and another one in the computations of intersections of the components of a Springer fiber.

Key words and phrases. Borel adjoint orbits; involutions and link patterns; Springer fibers and orbital varieties; Young tableaux.

This work was partially supported by the EEC program RTN-grant "Liegrits".

1.2. Let us first recall in short Jordan-Gerstenhaber theory.

Let $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0) \vdash n$ be a partition of n (ordered in decreasing order). Set $\lambda^* := \{\lambda_1^* \geq \lambda_2^* \geq \dots \geq \lambda_l^* > 0\}$ to be the conjugate partition, that is $\lambda_i^* = \#\{j \mid \lambda_j \geq i\}$.

The unique eigenvalue of a nilpotent matrix is 0, so that Jordan form of $u \in \mathfrak{N}$ is defined completely by the lengths of its Jordan blocks, which is defined in turn by some partition of n . Since each nilpotent orbit has a unique Jordan form up to the order of Jordan blocks we have a bijection between nilpotent orbits and ordered partitions. We can write $J(u) := \lambda$ and $\mathcal{O}_\lambda := \mathcal{O}_u$ where λ is the corresponding partition.

Let $\overline{V}$ denote the closure of variety V with respect to Zariski topology. The geometry of $\overline{\mathcal{O}_\lambda}$ is described combinatorially in terms of partitions. We begin with the formula of the dimension of an orbit:

$$\dim \mathcal{O}_\lambda = n^2 - \sum_{i=1}^l (\lambda_i^*)^2. \quad (*)$$

Our next goal is a combinatorial description of the closure of an orbit.

A partial order on partitions (called a dominance order) is defined as follows. Let $\lambda, \mu \vdash n$ be ordered in decreasing order $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \lambda_k > 0)$ and $\mu = (\mu_1 \geq \mu_2 \geq \dots \mu_l > 0)$. Set $\lambda \geq \mu$ if for any $j : 1 \leq j \leq \min(k, l)$ one has

$$\sum_{i=1}^j \lambda_i \geq \sum_{i=1}^j \mu_i.$$

Let $\coprod$ denote a disjoint union. By a theorem of M. Gerstehaber (cf. [4] for example)

$$\overline{\mathcal{O}_\lambda} = \coprod_{\mu \leq \lambda} \mathcal{O}_\mu.$$

Given a partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_k > 0)$ we define the corresponding Young diagram D_λ of λ to be an array of k rows of cells starting on the left with the i -th row containing λ_i cells.

Young diagrams are a very convenient visualization of the partitions with respect to nilpotent orbits. Indeed, the dual partition used in $(*)$ is simply the list of lengths of the columns of the corresponding Young diagrams. Also, $\mu < \lambda$ if D_μ is obtained from D_λ by pushing down some cells. In particular, the cover of a given partition (that is all the maximal partitions strictly smaller than the given one) with respect to the dominance order is described easily in the terms of Young diagrams.

Indeed, let $\mu = (\mu_1, \dots)$ be in the cover of $\lambda = (\lambda_1, \dots)$. Then μ is obtained from λ in one of two ways:

- (i) There exists i such that $\lambda_i - \lambda_{i+1} \geq 2$. Then $\mu_j = \lambda_j$ for $j \neq i, i+1$ and

$$\mu_i = \lambda_i - 1, \quad \mu_{i+1} = \lambda_{i+1} + 1.$$

In this case $\dim \mathcal{O}_\mu = \dim \mathcal{O}_\lambda - 2$.

- (ii) There exists i such that $\lambda_{i+1} = \lambda_{i+2} = \cdots = \lambda_{i+k} = \lambda_i - 1$ for some $k \geq 1$ and $\lambda_{i+k+1} = \lambda_i - 2$. Then $\mu_j = \lambda_j$ for $j \neq i, i+k+1$ and

$$\mu_i = \lambda_i - 1, \quad \mu_{i+k+1} = \lambda_i - 1.$$

In this case $\dim \mathcal{O}_\mu = \dim \mathcal{O}_\lambda - 2(k+1)$.

In particular the cover of a nilpotent orbit is not equidimensional in general.

The above result can be described by Young diagrams as follows.

In the first case D_μ is obtained from D_λ by pushing one box down one row (and possible across several columns). For example,

$$D_\lambda = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & X \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array}, \quad D_\mu = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & X & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array}.$$

In the second case diagram D_μ is obtained from D_λ by pushing one box across one column (and possible down several rows). For example,

$$D_\lambda = \begin{array}{|c|c|c|c|} \hline & & & X \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}, \quad D_\mu = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & X & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}.$$

1.3. In the paper we show that link patterns play the same role for the combinatorial description of geometry of **B** orbits in $\mathcal{X}$ as Young diagrams for the description of geometry of nilpotent orbits. First we describe the bijection between **B** orbits in $\mathcal{X}$ and link patterns.

Let $\mathbf{S}_n^2 := \{\sigma \in \mathbf{S}_n \mid \sigma^2 = Id\}$ be the subset of involutions of the symmetric group $\mathbf{S}_n$. For any $\sigma \in \mathbf{S}_n^2$ let N_σ be the matrix obtained from the representation matrix of σ by erasing the lower-triangular part (including the main diagonal), that is


$$(N_\sigma)_{i,j} := \begin{cases} 1 & \text{if } i < j \text{ and } \sigma(i) = j; \\ 0 & \text{otherwise.} \end{cases}$$

One can see at once that $N_\sigma \in \mathcal{X}$ for any $\sigma \in \mathbf{S}_n^2$. Moreover by [9, 2.2] one has

$$\mathcal{X} = \coprod_{\sigma \in \mathbf{S}_n^2} \mathcal{B}_{N_\sigma}.$$

Thus, $\mathbf{B}$ orbits in $\mathcal{X}$ are labelled by involutions of $\mathbf{S}_n$. Put $\mathcal{B}_\sigma := \mathcal{B}_{N_\sigma}$.

Given $\sigma \in \mathbf{S}_n^2$ the corresponding link pattern P_σ is an array of n points on a (horizontal) line where points $i \neq j$ are connected by an arc if $\sigma(i) = j$. We will call such points end points of an arc. A point i satisfying $\sigma(i) = i$ is called a fixed point of P_σ . For example, for $\sigma = (1, 3)(2, 6)(4, 7) \in \mathbf{S}_7^2$ one has



The only fixed point of P_σ is 5.

Remark. I found the notion of link pattern in the papers in combinatorics and mathematical physics [19], [6], [1]. In general link patterns considered there are patterns with maximal possible number of arcs and a link pattern is a circle with n points rather than a line. Respectively the authors use a word "arch" not "arc". I adopted the notion of a link pattern from them, however it seems to me that the word "arc" is more appropriate in the representation given here.

1.4. Let us explain the role of link patterns in the description of geometry of $\mathbf{B}$ orbits in $\mathcal{X}$.

Let $\mathcal{B}_\sigma$ be an orbit in $\mathcal{X}$ and P_σ be the corresponding link pattern. We begin with formula of $\dim \mathcal{B}_\sigma$ in terms of a link pattern. Given $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$ (in cyclic form).

- Put $\ell(P_\sigma) := k$ to be the number of arcs in P_σ (we will call it the length of a pattern);
- Arcs $(i, j), (i', j')$ where $i < i'$ of P_σ are called intersecting if $i < i' < j$ and $j' > j$. We say that arcs $(i, j), (i', j')$ have a crossing in this case. Put $c(P_\sigma)$ to be the number of crossings of arcs in P_σ ;
- Let $\{p_s\}_{s=1}^{n-2k}$ be the set of fixed points of P_σ . For $s : 1 \leq s \leq n - 2k$ put $f_{p_s}(P_\sigma)$ to be the number of arcs (i_t, j_t) over p_s (that is such that $i_t < p_s < j_t$) and $f(P_\sigma) := \sum_{s=1}^{n-2k} f_{p_s}(P_\sigma)$.

In the example above $\ell(P_\sigma) = 3$, $c(P_\sigma) = 2$, and $f(P_\sigma) = f_5(P_\sigma) = 2$.

As we show in 3.1 one has

$$\dim \mathcal{B}_\sigma = \ell(P_\sigma) \cdot (n - \ell(P_\sigma)) - c(P_\sigma) - f(P_\sigma).$$

Thus, in our example $\dim \mathcal{B}_\sigma = 3 \cdot 4 - 2 - 2 = 8$.

Note that $\mathbf{S}_n^2$ can be stratified by the length of link patterns. Put $\mathbf{S}_n^2(k) := \{\sigma \in \mathbf{S}_n^2 \mid \ell(P_\sigma) = k\}$. One has $\mathbf{S}_n^2 = \coprod_{k \leq \frac{n}{2}} \mathbf{S}_n^2(k)$.

Note that this stratification partitions $\mathbf{B}$ orbits in $\mathcal{X}$ into sets belonging to the same nilpotent orbit. Indeed, put $\mathcal{O}_\sigma := \mathcal{O}_{N_\sigma}$, then $\mathcal{O}_\sigma = \mathcal{O}_{\sigma'}$ iff $\ell(P_\sigma) = \ell(P_{\sigma'})$. This is because a nilpotent orbit of nilpotent order 2 is completely defined by the rank of its representing matrix and $\text{Rank}(N_\sigma) = \ell(P_\sigma)$.

1.5. Further we describe $\overline{\mathcal{B}}_\sigma$ in terms of P_σ .

For $i, j : 1 \leq i \leq j \leq n$ let $[i, j] := \{s\}_{s=i}^j$ be the set of integer points of a segment $[i, j]$. For an arc $(i_s, j_s) \in P_\sigma$ we put $(i_s, j_s) \in [i, j]$ if $i \leq i_s, j_s \leq j$. For $i, j : 1 \leq i < j \leq n$ let $(R_\sigma)_{i,j}$ be the number of arcs in P_σ belonging to $[i, j]$. Just for convenience we can represent R_σ as an $n \times n$ matrix with integer non negative entries by setting $(R_\sigma)_{i,j} = 0$ for any $i \geq j$. In our example

$$R_\sigma = \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Define a partial order on $\mathbf{M}_n(\mathbb{R})$ by $A \succeq B$ if for any $i, j : 1 \leq i, j \leq n$ one has $(A)_{i,j} \geq (B)_{i,j}$. This order induces a partial order on involutions by putting $\sigma \succeq \sigma'$ if $R_\sigma \succeq R_{\sigma'}$. Then, by 2.4 and 3.2 one has

$$\overline{\mathcal{B}}_\sigma = \coprod_{\sigma' \preceq \sigma} \mathcal{B}_{\sigma'}.$$

1.6. Let us apply the previous result to the cover of a given σ with respect to order $\preceq$. We describe the cover in terms of link patterns only.

First, notice that nilpotent orbits of nilpotent order 2 are ordered linearly with respect to a dominance order. Indeed, given λ, μ corresponding to nilpotent orbits of nilpotent order 2 then $\lambda^* = (n - k, k)$ and $\mu^* = (n - m, m)$. One has $\mu \leq \lambda$ iff $m \leq k$ otherwise $\mu > \lambda$. Thus, for any nilpotent orbits $\mathcal{O}, \mathcal{O}'$ of nilpotent order 2 one has either $\overline{\mathcal{O}}' \subset \overline{\mathcal{O}}$ or $\overline{\mathcal{O}}' \supset \overline{\mathcal{O}}$. In particular the cover of partition $(n - k, k)^*$ consists of the unique partition $(n - k + 1, k - 1)^*$ (it is of type (ii) described in 1.2).

Let $\mathcal{C}(\sigma)$ denote the cover for a given $\sigma \in \mathbf{S}_n^2$ with respect to order $\preceq$. In 3.9 we show that $\sigma' \in \mathcal{C}(\sigma)$ iff $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$ so that the cover with respect to order $\preceq$ is equidimensional.

However, from geometric point of view there are two interesting subsets of elements smaller than σ . The first one is $\mathcal{N}(\sigma)$ – the subset of maximal elements smaller than σ in the set of patterns of smaller length. The second one is $\mathcal{D}(\sigma) := \mathcal{C}(\sigma) \cap \mathbf{S}_n^2(\ell(P_\sigma))$.

Let $\sigma \in \mathbf{S}_n^2(k)$. Geometric meaning of $\mathcal{N}(\sigma)$ is provided by

$$\overline{\mathcal{B}}_\sigma \cap (\overline{\mathcal{O}}_\sigma \setminus \mathcal{O}_\sigma) = \overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{O}}_{(n-k+1, k-1)^*} = \bigcup_{\sigma' \in \mathcal{N}(\sigma)} \overline{\mathcal{B}}_{\sigma'}.$$

As for $\mathcal{D}(\sigma)$ it helps to understand the structure of $\overline{\mathcal{B}}_\sigma$ inside of $\mathcal{O}_\sigma$. In particular for $\sigma, \sigma' \in \mathbf{S}_n^2(k)$ such that $\dim \mathcal{B}_\sigma = \dim \mathcal{B}_{\sigma'}$ one has $\text{codim}_{\overline{\mathcal{B}}_\sigma}(\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'}) = 1$ if and only if $\mathcal{D}(\sigma) \cap \mathcal{D}(\sigma') \neq \emptyset$.

We give the combinatorial description in terms of link patterns of both $\mathcal{N}(\sigma)$ and $\mathcal{D}(\sigma)$. The description of $\mathcal{N}(\sigma)$ in terms of link patterns is very simple. We call an arc of P_σ **external** if there is no arc over it, that is $(i, j) \in P_\sigma$ is external if for any arc $(i', j') \neq (i, j)$ in P_σ one has $(i, j) \notin [i', j']$.

All the elements of $\mathcal{N}(\sigma)$ are obtained from σ by deleting an external arc. So the number of elements in $\mathcal{N}(\sigma)$ is equal to the number of external arcs in σ and all the elements in $\mathcal{N}(\sigma)$ are in $\mathbf{S}_n^2(\ell(P_\sigma) - 1)$.

Note that in general

- the set $\{\mathcal{B}_{\sigma'} \mid \sigma' \in \mathcal{N}(\sigma)\}$ is not equidimensional;
- $\mathcal{N}(\sigma)$ is **not** a subset of $\mathcal{C}(\sigma)$.

$\mathcal{D}(\sigma)$ consists of 3 possible types of elements. We describe them in 3.4-3.6.

1.7. All the results mentioned above are based on [11] and we use heavily the results of this paper in our proofs. The advantage of the representation of involutions by link patterns is that the complex combinatorics of [11] becomes easy and natural in this language.

The fact that $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$ for any $\sigma' \in \mathcal{C}(\sigma)$ was shown already in [14]. However, the proof provided there involves a lot of computations and here using the technique of link patterns we prove it in a few lines.

1.8. Let us use the technique of link patterns to study orbital varieties of nilpotent order 2 and components of Springer fibers of nilpotent order 2. To do this we consider link patterns without intersecting arcs and without fixed points under the arcs. They are the most simple link patterns from the combinatorial point of view on one hand and the most important objects for the applications to representation theory on the other hand.

By 1.4 $\dim \mathcal{B}_\sigma = (n - k)k$ (that is $\mathcal{B}_\sigma$ is of maximal possible dimension) iff P_σ is such link pattern. Note that by 1.2 $\dim \mathcal{O}_{(n-k, k)^*} = n^2 - k^2 - (n - k)^2 = 2k(n - k)$. Since in general $\dim \mathcal{B}_u \leq 0.5 \dim \mathcal{O}_u$ for $u \in \mathfrak{n}$ we get that $\mathcal{B}_\sigma$ corresponding to link patterns without intersecting arcs and without fixed points under the arcs are **B** orbits of maximal possible dimension. Moreover, their closures in $\mathcal{O}_\sigma$ are Lagrangian subvarieties of $\mathcal{O}_\sigma$, called orbital varieties.

Let us give a brief general description of an orbital variety (cf. 2.2 for the details). Let $\mathcal{O}$ be a nilpotent orbit in a semisimple Lie algebra $\mathfrak{g}$. Let us fix some triangular

decomposition $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \mathfrak{n}^-$. The irreducible components of $\mathcal{O} \cap \mathfrak{n}$ are called orbital varieties associated to $\mathcal{O}$. Note that $\mathcal{O}$ is a symplectic manifold and as it was shown by N. Spaltenstein, R. Steinberg and A. Joseph an orbital variety is a Lagrangian subvariety of $\mathcal{O}$. According to orbit method Lagrangian subvarieties should play an important role in the representation theory.

Let us explain the role of orbital varieties in $\mathfrak{g} = \mathfrak{sl}_n$ considered in the paper. In this case orbital varieties associated to $\mathcal{O}_\lambda$ are parameterized by standard Young tableaux corresponding to Young diagram D_λ . Let $\mathbf{Tab}_\lambda$ be the set of standard Young tableaux corresponding to D_λ . For $T \in \mathbf{Tab}_\lambda$ we denote by $\mathcal{V}_T$ the corresponding orbital variety associated to $\mathcal{O}_\lambda$.

By a result of A. Joseph primitive ideals of the enveloping algebra $U(\mathfrak{g})$ corresponding to a highest weight module with an integral weight are in bijection with standard Young tableaux. Let I_T be the primitive ideal corresponding to $T \in \mathbf{Tab}_\lambda$. As it is shown in [7] an orbital variety closure $\overline{\mathcal{V}}_T$ is the associated variety of I_T . As those they play a key role in the theory of primitive ideals. The details on the theory above can be found in [5].

Let S, T be Young tableaux with two columns. By [10] $I_T \subset I_S$ if and only if $\overline{\mathcal{V}}_T \supset \overline{\mathcal{V}}_S$. Further in [11] it was shown that for $T \in \mathbf{Tab}_{(n-k,k)^*}$ one has that the closure of $\mathcal{V}_T$ is a union of orbital varieties and the combinatorial description of this closure in terms of Young tableaux was obtained. However the combinatorial description in [11, 4.3] is too complex to be satisfactory. On the other hand, as it was noted already in [8] each orbital variety $\mathcal{V}_T$ of nilpotent order two admits the unique $\mathbf{B}$ orbit. Let us denote the corresponding involution by σ_T . In 4.2 we simplify this description using link patterns and get a simple answer in terms of $\mathcal{N}(\sigma_T)$.

1.9. As it was shown in [12] the bijection between orbital varieties associated to a nilpotent orbit $\mathcal{O}_\lambda$ and components of Springer fiber $\mathcal{F}_x$ where $x \in \mathcal{O}_\lambda$ is extended to the intersections: Let $S, T \in \mathbf{Tab}_\lambda$ and let $\mathcal{V}_T, \mathcal{V}_S$ be orbital varieties associated to $\mathcal{O}_\lambda$ and $\mathcal{F}_T, \mathcal{F}_S$ be the corresponding components of $\mathcal{F}_x$. The number of irreducible components of $\mathcal{V}_T \cap \mathcal{V}_S$ and their codimensions are equal to the number of irreducible components of $\mathcal{F}_T \cap \mathcal{F}_S$ and their codimensions. Via this bijection orbital varieties play a key role in Springer theory.

This bijection together with the theory of $\mathbf{B}$ orbits of nilpotent order 2 gave the opportunity to compute for the first time the intersections of the components of Springer fiber when these components are singular. These intersections were considered in [13], [14]. Here we use link patterns to simplify and extend these results. We start with a general theory of intersections of the closures of $\mathbf{B}$ orbits of nilpotent order 2 belonging to $\mathcal{O}_{(n-k,k)^*}$. Let $\sigma, \sigma' \in \mathbf{S}_n^2(k)$. The general picture of $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'} \cap \mathcal{O}_{(n-k,k)^*}$ was considered in [13], [14]. As it is shown there these intersections can be very complex. In particular, there are examples in these papers of a reducible intersection of codimension 1 and of a reducible intersection of higher codimension


which is not equidimensional (in 5.1 we give an example of an intersection of codimension 1 which is not equidimensional). To compute $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'} \cap \mathcal{O}_{(n-k,k)^*}$ we use matrix $R_{\sigma,\sigma'}$ defined by

$$(R_{\sigma,\sigma'})_{i,j} := \min\{(R_\sigma)_{i,j}, (R_{\sigma'})_{i,j}\}$$

As it was shown in [13, 5.6, 5.7] this intersection is non empty and it is a union of all $\mathcal{B}_{\sigma''}$ where $\sigma'' \in \mathbf{S}_n^2(k)$ are such that $R_{\sigma''} \preceq R_{\sigma,\sigma'}$. In particular this intersection is irreducible iff there exists $\sigma'' \in \mathbf{S}_n^2(k)$ such that $R_{\sigma,\sigma'} = R_{\sigma''}$.

To simplify the results on intersections we pass from link patterns to so called (generalized) meanders. To do this we draw two link patterns on the same line of points in such a way that the arcs of the first link pattern are drawn upward and the arcs of the second one are drawn downward.

For example, consider $\sigma = (1,3)(2,5)(4,7)$, $\sigma' = (1,4)(3,5)(6,7) \in \mathbf{S}_7^2$. Their generalized meander $M_{\sigma,\sigma'}$ is



These generalized meanders are useful for computing $R_{\sigma,\sigma'}$.

Taking a meander built of 2 link patterns corresponding to maximal $\mathbf{B}$ orbits in $\mathcal{O}_{(n-k,k)^*}$ we get a classical meander (that is without crossing arcs) without fixed points under any (upward or downward) arc. As it is shown in [14] for $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ if $\text{codim}_{\mathcal{V}_T}(\mathcal{V}_S \cap \mathcal{V}_T) = 1$ then $\mathcal{V}_S \cap \mathcal{V}_T$ is irreducible. The combinatorial description of pairs S, T with intersection of codimension 1 is provided there.

However, the consideration of these intersections in terms of meanders first of all simplifies the description and the proofs and further connects our results to Temperley-Lieb representations of Hecke algebras and Kazhdan-Lusztig data.

Let us formulate the result. A meander M_{σ_T, σ_S} is called even if any its connected part consists of even number of arcs. Otherwise M_{σ_T, σ_S} is called odd. The connected part consisting of even number of arcs may be either a closed path (we call it a loop) or an open path (we call it an even interval). As we show in 5.3 for $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ one has $\text{codim}_{\mathcal{V}_T}(\mathcal{V}_S \cap \mathcal{V}_T) = 1$ if and only if M_{σ_S, σ_T} is an even meander with $k-1$ loops. The irreducibility of such intersections is obvious from the combinatorics of the meanders.

B.W. Westbury in [18] computed the inner product matrix for the Kazhdan-Lusztig basis for a two row shape tableaux in terms of Temperley-Lieb cup diagrams. Note that for $T \in \mathbf{Tab}_{(n-k,k)}$ the corresponding cup diagram defined in [18] is exactly $P_{\sigma_{T^t}}$ where $T^t \in \mathbf{Tab}_{(n-k,k)^*}$ is a transposed tableau. Given $S, T \in \mathbf{Tab}_{(n-k,k)}$ let c_T, c_S be the corresponding Kazhdan-Lusztig basis vectors from [18]. As it is shown

in [18, 7] their inner product $\langle c_S, c_T \rangle$ is always either 0 or $(t + t^{-1})^r$ where $0 \leq r \leq k - 1$. Moreover, restating his results in the language of meanders we get $\langle c_S, c_T \rangle = (t + t^{-1})^r$ if and only if the corresponding meander M_{σ_S, σ_T} is an even meander with r loops.

F.Y.C. Fung in [2] computed the intersections of the components of a Springer fiber corresponding to a nilpotent element with two Jordan blocks, that is $\mathcal{F}_T \cap \mathcal{F}_S$ where $T, S \in \mathbf{Tab}_{(n-k, k)}$. In this case $\mathcal{F}_T$ is nonsingular and as he showed all the intersections are either irreducible or empty. Using the technique similar to those used in [18] he showed that $\mathcal{F}_S \cap \mathcal{F}_T = \emptyset$ if and only if $\langle c_T, c_S \rangle = 0$ and if $\langle c_T, c_S \rangle \neq 0$ then $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) = k - \deg \langle c_T, c_S \rangle$.

Consider $S, T \in \mathbf{Tab}_{(n-k, k)}$. Combining the above we get that $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) = 1$ if and only if $\text{codim}_{\mathcal{F}_{T^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t}) = 1$ and this happens iff $\langle c_S, c_T \rangle = (t + t^{-1})^{k-1}$.

However the pictures of the intersections of higher codimension in two-row and two-column cases are very different. Note that for $\sigma, \sigma' \in \mathbf{S}_n^2(k)$ one has $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'} \cap \mathcal{O}_{(n-k, k)^*} \neq \emptyset$, thus, in particular for an odd meander M_{σ_S, σ_T} one has $\langle c_{S^t}, c_{T^t} \rangle = 0$ and $\mathcal{F}_T \cap \mathcal{F}_S \neq \emptyset$. On the other hand, for even meanders with less than $k - 1$ loops it can occur that $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) > \text{codim}_{\mathcal{F}_{T^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t})$ as we show by example in 5.5. The general theory of intersections of higher codimension is too complex from combinatorial point of view and not that interesting from geometric point of view so we do not discuss it here.

Also, the intersections of higher codimension are almost always reducible. Again, the general theory is too involved combinatorially to explain it here so we only provide in 5.6 a very simple combinatorial sufficient condition for the reducibility of an intersection which shows that in general the intersections of higher codimension are reducible.

In the end of the paper one can find the index of notation in which symbols appearing frequently are given with the subsection where they are defined. I hope that this will help the reader to find his way through the paper.

Acknowledgements. I am very grateful to A. Elashvili for the discussions on biparabolic algebras and the role of meanders in their description. He was first to introduce the world of meanders to me. Through these discussions I realized the role that they (and link patterns) play in the description of $\mathbf{B}$ orbits of nilpotent order 2 and intersections of the components of a Springer fiber of nilpotent order 2. I am also very grateful to A. Joseph for many fruitful discussions during various stages of this work.

2. $\mathbf{B}$ ORBITS IN $\mathcal{X}$ AND INVOLUTIONS

2.1. Put $\mathfrak{B}_{(n-k,k)} = \coprod_{\sigma \in \mathbf{S}_n^2(k)} \mathcal{B}_\sigma$. As we noted in 1.4 $\mathfrak{B}_{n-k,k} = \mathcal{X} \cap \mathcal{O}_{(n-k,k)}^*$ so that $\mathcal{X} = \coprod_{k \leq \frac{n}{2}} \mathfrak{B}_{(n-k,k)}$.

We start with the computation of the dimension of $\mathcal{B}_\sigma$. All over this paper we write a cycle in round brackets (i, j) when the entries are ordered in the increasing order, i.e. $i < j$. We write it in double round brackets $((i, j))$ when the entries $i \neq j$ are not ordered. Write $\sigma \in \mathbf{S}_n^2(k)$ as a product of disjoint cycles of length 2, that is $\sigma = (i_1, j_1)(i_2, j_2) \dots (i_k, j_k)$ where $i_s < j_s$ and $\{i_s, j_s\} \cap \{i_t, j_t\} = \emptyset$ for any $1 \leq s \neq t \leq k$. We write $(i, j) \in \sigma$ or $(i, j) \in P_\sigma$ if $(i, j) = (i_s, j_s)$ for some $s : 1 \leq s \leq k$. For $(i, j) \in \sigma$ put

$$q_{(i,j)}(\sigma) := \#\{i_p < i \mid j_p < j\} + \#\{j_p \mid j_p < i\}. \quad (*)$$

Example. Take $\sigma = (1, 6)(3, 4)(5, 7) \in \mathbf{S}_7^2(3)$. Then $\ell(P_\sigma) = 3$ and $q_{(1,6)}(\sigma) = 0$, $q_{(3,4)}(\sigma) = 0$, $q_{(5,7)}(\sigma) = 2 + 1 = 3$.

By [9, 3.1] one has

Theorem. For $\sigma = (i_1, j_1)(i_2, j_2) \dots (i_k, j_k) \in \mathbf{S}_n^2(k)$ one has

$$\dim \mathcal{B}_\sigma = kn - \sum_{s=1}^k (j_s - i_s) - \sum_{s=1}^k q_{(i_s, j_s)}(\sigma).$$

2.2. What is the maximal dimension of an orbit in $\mathfrak{B}_{(n-k,k)}$? We would like to provide a detailed answer to this question. To do this we consider in detail orbital varieties defined briefly in 1.8.

Let $\mathbf{G}$ be a connected semisimple finite dimensional complex algebraic group and let $\mathfrak{g} = \text{Lie}(\mathbf{G})$ be the corresponding semisimple Lie algebra. Let $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \mathfrak{n}^-$ be its triangular decomposition. For $u \in \mathfrak{n}$ let $\mathcal{O}_u = \mathbf{G}.u$ be its adjoint (nilpotent) orbit. Note that by the identification of $\mathfrak{g}$ with $\mathfrak{g}^*$ through the Killing form $\mathcal{O}_u$ is provided a symplectic structure.

Consider $\mathcal{O}_u \cap \mathfrak{n}$. This variety is reducible and its irreducible components are called orbital varieties associated to $\mathcal{O}_u$. Let $\mathcal{V}$ be an orbital variety associated to $\mathcal{O}_u$. By [16] and [17] one has $\dim \mathcal{V} = \frac{1}{2} \dim \mathcal{O}_u$. (Moreover as it was pointed out in [5] this implies that an orbital variety is a Lagrangian subvariety of the nilpotent orbit it is associated to.)

In particular, if $\mathbf{G} = SL_n$ nilpotent orbits are described by Young diagrams as we explained in 1.2. In this case there exists a very nice combinatorial characterization of orbital varieties in terms of Young tableaux. Recall that a Young tableau T associated to Young diagram D_λ is obtained by filling the boxes of D_λ with numbers $1, \dots, n$ so that the numbers increase in rows from left to right and in columns from top to bottom. Given a Young tableau T associated to D_λ its shape is defined to be λ and denoted by $\text{sh}(T)$. Given $u \in \mathfrak{n} \cap \mathcal{O}_\lambda$ its Young diagram is again defined

to be λ and denoted by $D_n(u)$, or simply $D(u)$. Now consider canonical projections $\pi_{1,n-i} : \mathfrak{n}_n \rightarrow \mathfrak{n}_{n-i}$ acting on a matrix by deleting the last i columns and the last i rows. Set $D_{n-1}(u) := D(\pi_{1,n-1}(u))$, $D_{n-2}(u) := D(\pi_{1,n-2}(u))$, $\dots$, $D_1(u) := D(\pi_{1,1}(u))$ and $\phi(u) := (D_1(u), D_2(u), \dots, D_n(u))$. Put 1 into the unique box of $D_1(u)$. For any $i : 1 \leq i \leq n-1$ note that $D_{i+1}(u)$ differs from $D_i(u)$ by a single "corner" box. Put $i+1$ into this box. This gives a bijection from the set of the chains $(D_1(u), D_2(u), \dots, D_n(u))$ to the set of standard Young tableaux T of shape $D_n(u)$. In other words, we view a standard Young tableau as a chain of Young diagrams. For $u \in \mathfrak{n}$ put $\varphi(u) := T$ if T corresponds to $\phi(u)$ under this bijection. Set $\nu_T := \{u \in \mathfrak{n} \mid \varphi(u) = T\}$.

By Spaltenstein [15] orbital varieties associated to $\mathcal{O}_\lambda$ are parameterized by standard Young tableaux of shape λ as follows. Let $\{T_i\}$ be the set of Young tableaux of shape λ . Set $\mathcal{V}_{T_i} := \overline{\nu}_{T_i} \cap \mathcal{O}_\lambda$. Then $\{\mathcal{V}_{T_i}\}$ is the set of orbital varieties associated to $\mathcal{O}_\lambda$.

Note that in particular by this construction $\dim \mathcal{B}_u \leq \frac{1}{2} \dim \mathcal{O}_u$. In general this inequality is strict. Moreover, obviously if $\dim \mathcal{B}_u = \frac{1}{2} \dim \mathcal{O}_u$ then $\mathcal{B}_u$ is dense in the corresponding orbital variety. Unfortunately, for $n \geq 6$ the vast majority of orbital varieties do not admit a dense **B** orbit. However, orbital varieties of nilpotent order two always admit a unique dense orbit. That gives the answer to the question posed above.

To formulate this answer let us consider the corresponding Young tableaux in detail. Let $\mathbf{Tab}_{(n-k,k)^*}$ be the set of Young tableaux of shape $(n-k, k)^*$. For $T \in$

$\mathbf{Tab}_{(n-k,k)^*}$ set $T = (T_1, T_2)$, where $T_1 = \begin{pmatrix} a_1 \\ \vdots \\ a_{n-k} \end{pmatrix}$ is the first column of T and

$T_2 = \begin{pmatrix} j_1 \\ \vdots \\ j_k \end{pmatrix}$ is the second column of T . It is enough to define the columns as sets

since the entries increase from top to bottom in the columns. We denote a column by $\langle T_i \rangle$ when we consider it as a set.

Put $\sigma_T = (i_1, j_1) \dots (i_k, j_k)$ where $i_1 = j_1 - 1$, and $i_s = \max\{d \in \langle T_1 \rangle \setminus \{i_1, \dots, i_{s-1}\} \mid d < j_s\}$ for any $s > 1$. For example, take

$$T = \begin{array}{cc} 1 & 4 \\ 2 & 5 \\ 3 & 7 \\ & 6 & 8 \end{array}$$

Then $\sigma_T = (3, 4)(2, 5)(6, 7)(1, 8)$.

Put $\mathcal{B}_T := \mathcal{B}_{\sigma_T}$. As it was shown in [8, 4.13]

Proposition. For $T \in \mathbf{Tab}_n^2$ one has $\overline{\mathcal{V}}_T = \overline{\mathcal{B}}_T$.

By [13, Remark 5.5] for $\sigma \in \mathbf{S}_n^2(k)$ one has $\dim \mathcal{B}_\sigma \leq k(n-k)$ and the equality is satisfied iff $\sigma = \sigma_T$ for some $T \in \mathbf{Tab}_{(n-k,k)^*}$. That is the number of $\mathcal{B}_\sigma \in \mathfrak{B}_{(n-k,k)}$ of dimension $k(n-k)$ is equal to the number of Young tableaux in $\mathbf{Tab}_{(n-k,k)^*}$ which is $\frac{n!}{k!(n-k)!} \frac{n-2k+1}{n-k+1}$.

Let $\dim \mathcal{B}_\sigma = k(n-k)$ where $\sigma = (i_1, j_1) \dots (i_k, j_k)$ then $\sigma = \sigma_T$ where $\langle T_2 \rangle = \{j_1, \dots, j_k\}$ and respectively $\langle T_1 \rangle = \{i\}_{i=1}^n \setminus \langle T_2 \rangle$.

2.3. In [11] the combinatorial description of $\overline{\mathcal{B}}_\sigma$ (with respect to Zariski topology) for $\sigma \in \mathbf{S}_n^2$ is provided. Let us formulate this result.

For $1 \leq i < j \leq n$ consider the canonical projections $\pi_{i,j} : \mathbf{n}_n \rightarrow \mathbf{n}_{j-i+1}$ acting on a matrix by deleting the first $i-1$ columns and rows and the last $n-j$ columns and rows. Define the rank matrix R_u of $u \in \mathbf{n}$ to be

$$(R_u)_{i,j} = \begin{cases} \text{Rank}(\pi_{i,j}(u)) & \text{if } i < j; \\ 0 & \text{otherwise.} \end{cases}$$

Obviously for any $y \in \mathcal{B}_u$ one has $R_y = R_u$ so that we can define $R_{\mathcal{B}_u} := R_u$.

Put $R_\sigma := R_{\mathcal{B}_\sigma} = R_{N_\sigma}$. Note that $\text{Rank}(\pi_{i,j}(N_\sigma))$ is simply the number of ones in matrix $\pi_{i,j}(N_\sigma)$ since all ones belong to different rows and columns.

Let $\mathbb{Z}^+$ be the set of non-negative integers. Put $\mathbf{R}_n^2 := \{R_\sigma \mid \sigma \in \mathbf{S}_n^2\}$. By [11, 3.1, 3.3] one has

Proposition. $R \in M_{n \times n}(\mathbb{Z}^+)$ belongs to $\mathbf{R}_n^2$ if and only if it satisfies

- (i) $R_{i,j} = 0$ if $i \geq j$;
- (ii) For $i < j$ one has $R_{i+1,j} \leq R_{i,j} \leq R_{i+1,j} + 1$ and $R_{i,j-1} \leq R_{i,j} \leq R_{i,j-1} + 1$;
- (iii) If $R_{i,j} = R_{i+1,j} + 1 = R_{i,j-1} + 1 = R_{i+1,j-1} + 1$ then
 - (a) $R_{i,k} = R_{i+1,k}$ for any $k < j$ and $R_{i,k} = R_{i+1,k} + 1$ for any $k \geq j$;
 - (b) $R_{k,j} = R_{k,j-1}$ for any $k > i$ and $R_{k,j} = R_{k,j-1} + 1$ for any $k \leq i$;
 - (c) $R_{j,k} = R_{j+1,k}$ and $R_{k,i} = R_{k,i-1}$ for any $k : 1 \leq k \leq n$.

Note that we respectively can define $\pi_{i,j} : \mathbf{Tab}_n \rightarrow \mathbf{Tab}_{j-i+1}$ by so called "jeu de taquin" (cf. [11], for example). Here we need only $\pi_{1,k} : \mathbf{Tab}_n \rightarrow \mathbf{Tab}_k$ where $\pi_{1,k}(T)$ is obtained by erasing boxes with $n, n-1, \dots, k+1$. As it is shown in [11] and one can easily see $\pi_{i,j}\mathcal{B}_T = \mathcal{B}_{\pi_{i,j}(T)}$.

2.4. Recall a partial order $\preceq$ on $\mathbf{M}_n(\mathbb{R})$ defined in 1.5. Its restriction to $\mathbf{R}_n^2$ defines a partial order $\preceq$ on $\mathbf{S}_n^2$ by putting $\sigma' \preceq \sigma$ if $R_{\sigma'} \preceq R_\sigma$. As it is shown in [11, 3.5]

Theorem. For any $\sigma \in \mathbf{S}_n^2$ one has

$$\overline{\mathcal{B}}_\sigma = \coprod_{\sigma' \preceq \sigma} \mathcal{B}_{\sigma'}.$$

2.5. Consider $\pi_{i,j}(\sigma)$ as an element of $\mathbf{S}_n^2$. Note that $(R_{\pi_{i,j}(\sigma)})_{s,t} = (R_\sigma)_{\max(i,s), \min(j,t)}$ which gives us a very easy lemma that we need in what follows.

Lemma. *For any $\sigma, \sigma' \in \mathbf{S}_n^2$ such that $\sigma \preceq \sigma'$ and for any $i, j : 1 \leq i < j \leq n$ one has*

- (i) $\pi_{i,j}(\sigma) \preceq \pi_{i,j}(\sigma')$
- (ii) $\pi_{i,j}(\sigma) \preceq \sigma$

2.6. As it is shown in [13, 5.6] there is a unique minimal element $\sigma_o(k)$ in $\mathbf{S}_n^2(k)$ (that is $\sigma \succeq \sigma_o(k)$ for any $\sigma \in \mathbf{S}_n^2(k)$). In particular for any $S' \subseteq \mathbf{S}_n^2(k)$ one has

$$\bigcap_{\sigma \in S'} \overline{\mathcal{B}}_\sigma \bigcap \mathcal{O}_{(n-k,k)^*} \neq \emptyset. \quad (*)$$

To find $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'}$ for $\sigma, \sigma' \in \mathbf{S}_n^2$ we use an intersection matrix $R_{\sigma, \sigma'}$ for $\sigma, \sigma' \in \mathbf{S}_n^2$ defined in 1.9. As it is shown in [13, 5.7],

Theorem. *For any $\sigma, \sigma' \in \mathbf{S}_n^2$ one has*

$$\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'} = \coprod_{R_{\sigma''} \preceq R_{\sigma, \sigma'}} \mathcal{B}_{\sigma''}$$

In particular, $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'}$ is irreducible if and only if $R_{\sigma, \sigma'} \in \mathbf{R}_n^2$. In that case $\overline{\mathcal{B}}_\sigma \cap \overline{\mathcal{B}}_{\sigma'} = \overline{\mathcal{B}}_\tau$ where $R_\tau = R_{\sigma, \sigma'}$.

3. $\mathcal{B}_\sigma$ IN TERMS OF LINK PATTERNS

3.1. In this section we translate all the results of the previous section into the language of link patterns. Recall $\ell(P_\sigma)$, $c(P_\sigma)$, $f(P_\sigma)$ from 1.4 and $[i, j]$ from 1.5. We put $[i, j] := \emptyset$ if $i > j$. For any point $m : 1 \leq m \leq n$ put $\mathbf{over}_m(P_\sigma)$ to be the set of arcs over m , that is $\mathbf{over}_m(P_\sigma) := \{(i, j) \in P_\sigma \mid m \in [i+1, j-1]\}$. For any $(i, j) \in P_\sigma$ put $\mathbf{over}_{(i,j)}(P_\sigma)$ to be the set of arcs over arc (i, j) , that is $\mathbf{over}_{(i,j)}(P_\sigma) := \{(i', j') \in P_\sigma \mid (i, j) \in [i'+1, j'-1]\}$

Theorem. *For $\sigma \in \mathbf{S}_n^2$ one has*

$$\dim \mathcal{B}_\sigma = \ell(P_\sigma) \cdot (n - \ell(P_\sigma)) - c(P_\sigma) - f(P_\sigma).$$

Proof. Our proof is based on the recalculating the expression of Theorem 2.1 in the language of link patterns.

Let $\sigma = (i_1, j_1) \dots (i_k, j_k)$. For a given arc $(i, j) \in P_\sigma$ we define

- $l_{(i,j)}(P_\sigma)$ to be the set arcs to the left of (i, j) , that is $l_{(i,j)}(P_\sigma) := \{(i', j') \in P_\sigma \mid (i', j') \in [1, i-1]\}$;
- $r_{(i,j)}(P_\sigma)$ to be the set of arcs to the right of (i, j) , that is $r_{(i,j)}(P_\sigma) := \{(i', j') \in P_\sigma \mid (i', j') \in [j+1, n]\}$.
- $\mathbf{under}_{(i,j)}(P_\sigma)$ to be the set of arcs under (i, j) that is $\mathbf{under}_{(i,j)}(P_\sigma) := \{(i', j') \in P_\sigma \mid (i', j') \in [i+1, j-1]\}$.

- $c_{(i,j)}^l(P_\sigma)$ to be the set of arcs intersecting with (i,j) on the left that is $c_{(i,j)}^l(P_\sigma) := \{(i', j') \in P_\sigma \mid i' \in [1, i-1] \text{ and } j' \in [i+1, j-1]\}$.
- $c_{(i,j)}^r(P_\sigma)$ to be the set of arcs intersecting with (i,j) on the right that is $c_{(i,j)}^r(P_\sigma) := \{(i', j') \in P_\sigma \mid i' \in [i+1, j-1] \text{ and } j' \in [j+1, n]\}$.
- recall that $\mathbf{over}_{(i,j)}(P_\sigma) := \{(i', j') \in P_\sigma \mid (i, j) \in [i'+1, j'-1]\}$.

Put $|S|$ to be cardinality of set S . Let us provide a few identities on the summing of the cardinalities of the sets defined above.

- (i) Note that the list of six types of arcs above includes all the possible positions of an arc $(i', j') \neq (i, j)$ of P_σ with respect to (i, j) . Thus, for any $(i, j) \in P_\sigma$ one has

$$|l_{(i,j)}(P_\sigma)| + |r_{(i,j)}(P_\sigma)| + |\mathbf{under}_{(i,j)}(P_\sigma)| + |\mathbf{over}_{(i,j)}(P_\sigma)| + |c_{(i,j)}^l(P_\sigma)| + |c_{(i,j)}^r(P_\sigma)| = k-1.$$

- (ii) Let $\{p_s\}_{s=1}^{n-2k}$ be the set of fixed points of P_σ . Put $f'_{(i,j)}(P_\sigma)$ to be the number of fixed points under the arc (i, j) then obviously $\sum_{s=1}^k f'_{(i_s, j_s)}(P_\sigma) = \sum_{s=1}^{n-2k} f_{p_s}(P_\sigma) = f(P_\sigma)$. Note also that for $(i, j) \in \sigma$ any $p \in [i+1, j-1]$ is either a fixed point, or some end point of $(i', j') \in \mathbf{under}_{(i,j)}(P_\sigma)$ or some end point of an arc intersecting (i, j) . Thus, one has

$$j - i - 1 = f'_{(i,j)}(P_\sigma) + 2|\mathbf{under}_{(i,j)}(P_\sigma)| + |c_{(i,j)}^l(P_\sigma)| + |c_{(i,j)}^r(P_\sigma)|.$$

- (iii) Note that $(i, j) \in c_{(i', j')}^l(P_\sigma)$ iff $(i', j') \in c_{(i,j)}^r(P_\sigma)$ so that

$$\sum_{s=1}^k |c_{(i_s, j_s)}^l(P_\sigma)| = \sum_{s=1}^k |c_{(i_s, j_s)}^r(P_\sigma)| = c(P_\sigma).$$

- (iv) Note also that arc $(i', j') \in l_{(i,j)}(P_\sigma)$ iff $(i, j) \in r_{(i', j')}(P_\sigma)$ so that

$$\sum_{s=1}^k |l_{(i_s, j_s)}(P_\sigma)| = \sum_{s=1}^k |r_{(i_s, j_s)}(P_\sigma)|.$$

- (v) Exactly in the same way $(i', j') \in \mathbf{under}_{(i,j)}(P_\sigma)$ iff $(i, j) \in \mathbf{over}_{(i', j')}(P_\sigma)$ so that

$$\sum_{s=1}^k |\mathbf{under}_{(i_s, j_s)}(P_\sigma)| = \sum_{s=1}^k |\mathbf{over}_{(i_s, j_s)}(P_\sigma)|.$$

- (vi) Finally note that $\#\{j_p \mid j_p < i\}$ (from $(*)$ of 2.1) is exactly $|l_{(i,j)}(P_\sigma)|$ and $\#\{i_p < i \mid j_p < j\}$ (from $(*)$ of 2.1) is $|l_{(i,j)}(P_\sigma)| + |c_{(i,j)}^l(P_\sigma)|$ so that

$$q_{(i,j)}(\sigma) = 2|l_{(i,j)}(P_\sigma)| + |c_{(i,j)}^l(P_\sigma)|.$$

In the following computations we omit the notation (P_σ) in all the sets defined above since we consider only P_σ here. Starting from Theorem 2.1 we get

$$\begin{aligned}
\dim \mathcal{B}_\sigma &= kn - \sum_{s=1}^k (j_s - i_s) - \sum_{s=1}^k q_{(i_s, j_s)}(\sigma) \quad \text{by (ii), (vi)} \\
&= kn - \sum_{s=1}^k (1 + f'_{(i_s, j_s)}(\sigma) + 2|\mathbf{under}_{(i_s, j_s)}| + |c_{(i_s, j_s)}^l| + |c_{(i_s, j_s)}^r|) \\
&\quad - \sum_{s=1}^k (2|l_{(i_s, j_s)}| + |c_{(i_s, j_s)}^l|) \quad \text{by (iii) and def. } f(\sigma) \\
&= nk - k - f(\sigma) - 3c(P_\sigma) - \sum_{s=1}^k (2|\mathbf{under}_{(i_s, j_s)}| + 2|l_{(i_s, j_s)}|) \quad \text{by (iv), (v)} \\
&= nk - k - f(\sigma) - 3c(P_\sigma) - \sum_{s=1}^k (|\mathbf{under}_{(i_s, j_s)}| + |\mathbf{over}_{(i_s, j_s)}| + |l_{(i_s, j_s)}| + |r_{(i_s, j_s)}|) \quad \text{by (i)} \\
&= nk - k - f(\sigma) - 3c(P_\sigma) - \sum_{s=1}^k (k - 1 - (|c_{(i_s, j_s)}^l| + |c_{(i_s, j_s)}^r|)) \quad \text{by (iii)} \\
&= nk - k - f(\sigma) - 3c(P_\sigma) - k(k - 1) + 2c(P_\sigma) = \\
&= k(n - k) - f(P_\sigma) - c(P_\sigma).
\end{aligned}$$

□

3.2. Let us translate the definitions of 2.3 into the language of link patterns.

For $i, j : 1 \leq i < j \leq n$ we define $\pi_{i,j}(P_\sigma)$ to be a link pattern from i to j . It consists only of arcs $(i', j') \in P_\sigma$ such that $(i', j') \in [i, j]$. On the other hand, there is 1 in place (i', j') of matrix N_σ where $i' \geq i$ and $j' \leq j$ iff there exists $(i', j') \in \sigma$. Hence $\ell(\pi_{i,j}(P_\sigma))$ is equal to number of ones in $\pi_{i,j}(N_\sigma)$ which in turn is $\text{Rank}(\pi_{i,j}(N_\sigma))$ as it was noted in 2.3. Thus,


$$(R_\sigma)_{i,j} = \begin{cases} \ell(\pi_{i,j}(P_\sigma)) & \text{if } i < j; \\ 0 & \text{otherwise;} \end{cases}$$

exactly as we have defined in 1.5. Together with 2.4 this provides the result announced in 1.5.

3.3. Now we can describe the cover of σ for our partial order in terms of link patterns. We start with the description of $\mathcal{N}(\sigma)$ defined in 1.6.

Recall from 1.6 that $(i, j) \in P_\sigma$ is called external if $\mathbf{over}_{(i,j)}(P_\sigma) = \emptyset$. Let $E(\sigma)$ be the set of external arcs in P_σ .

For example, let $\sigma = (1, 3)(2, 7)(4, 5)$ then P_σ is



and $E(\sigma) = \{(1, 3), (2, 7)\}$.

For $(i, j) \in \sigma$ put $\sigma_{(i,j)}^-$ to be an involution obtained from σ by omitting (i, j) . Respectively $P_{\sigma_{(i,j)}^-}$ is obtained from P_σ by erasing arc (i, j) .

By [11, 3.7, 3.9] one has

Proposition. $\mathcal{N}(\sigma) = \{\sigma_{(i,j)}^- \mid (i, j) \in E(\sigma)\}$

3.4. $\mathcal{D}(\sigma)$ defined in 1.6 is more involved. Its description given in [11, 3.11-3.14] is rather complicated. However it is much more short and elegant in the language of link patterns. $\mathcal{D}(\sigma)$ contains of three types of elements which we describe in three following subsections. The main idea which is very clear on the level of link patterns is that the elements of $\mathcal{D}(\sigma)$ are obtained either by moving a left (resp. right) end point of an arc to the nearest fixed point to the left (resp. to the right) of it, so that we get a new fixed point under the arc, or by crossing two arcs. If $P_{\sigma'}$ is obtained from P_σ in such way, then $\mathcal{B}_{\sigma'}$ must be of codimension 1 in $\overline{\mathcal{B}}_\sigma$. A more delicate point is that all the elements of $\mathcal{D}(\sigma)$ are obtained in such way. This is exactly the content of Proposition 3.15 of [11].

We begin with constructing of a link pattern obtained from a given one by moving one end point of an arc to the nearest fixed point.

Given $\sigma = (i_1, j_1) \dots (i_k, j_k)$ we put $\langle \sigma \rangle := \langle P_\sigma \rangle := \{i_1, j_1, \dots, i_k, j_k\}$ to be the set of end points of P_σ .

For $i \in \langle \sigma \rangle$ and $f \notin \langle \sigma \rangle$ put $\sigma_{i \rightarrow f}$ to be an involution obtained from σ by changing an arc $((i, j))$ to $((f, j))$. We write the arcs in double round brackets since the definition is not connected to the ordering of the ends of an arc.

For $(i, j) \in \sigma$ if there exist fixed points to the left of arc (i, j) let m be the largest among them, that is $m = \max\{p < i \mid p \notin \langle \sigma \rangle\}$. If in addition $\mathbf{over}_{(i,j)}(P_\sigma) \cap [m + 1, n] = \emptyset$ (or in other words $\mathbf{over}_{(i,j)}(P_\sigma) \subseteq \mathbf{over}_m(P_\sigma)$) we put $\sigma_{\curvearrowright(i,j)} := \sigma_{i \rightarrow m}$. Otherwise put $\sigma_{\curvearrowright(i,j)} = \emptyset$.

Let $\sigma_{\curvearrowright(i,j)} = \sigma_{i \rightarrow m}$. Note that by 3.2 we get


$$(R_{\sigma_{\curvearrowright(i,j)}})_{s,t} = \begin{cases} (R_\sigma)_{s,t} - 1 & \text{if } m < s \leq i \text{ and } t \geq j; \\ (R_\sigma)_{s,t} & \text{otherwise.} \end{cases}$$

Thus, if $\sigma_{\curvearrowright(i,j)} \neq \emptyset$ then $\sigma_{\curvearrowright(i,j)} \prec \sigma$.


Exactly in the same way, for $(i, j) \in \sigma$ if there exist fixed points to the right of (i, j) let m be the minimal among them, that is $m = \min\{p > j \mid p \notin \langle \sigma \rangle\}$. If in addition $\mathbf{over}_{(i,j)}(P_\sigma) \cap [1, m - 1] = \emptyset$ (that is again $\mathbf{over}_{(i,j)}(P_\sigma) \subseteq \mathbf{over}_m(P_\sigma)$) we put $\sigma_{(i,j) \curvearrowright} := \sigma_{j \rightarrow m}$. Otherwise put $\sigma_{(i,j) \curvearrowright} = \emptyset$.

And again, exactly as in case $\sigma_{\curvearrowright(i,j)}$ if $\sigma_{(i,j) \curvearrowright} \neq \emptyset$ then $\sigma_{(i,j) \curvearrowright} \prec \sigma$.


For example, let $\sigma = (1, 6)(3, 5)(4, 7) \in \mathbf{S}_8^2$.




Then $\sigma_{\curvearrowright(1,6)} = \emptyset$ and $\sigma_{(1,6)\curvearrowright} = (1,8)(3,5)(4,7)$ so that



$\sigma_{\curvearrowright(3,5)} = (1,6)(2,5)(4,7)$ and $\sigma_{(3,5)\curvearrowright} = \emptyset$ so that



Finally, $\sigma_{\curvearrowright(4,7)} = (1,6)(3,5)(2,7)$ and $\sigma_{(4,7)\curvearrowright} = (1,6)(3,5)(4,8)$ so that



As an easy corollary of Theorem 3.1 we get

Proposition. *Let σ' be either $\sigma_{\curvearrowright(i,j)}$ or $\sigma_{(i,j)\curvearrowright}$. Then $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$.*

Proof. By our note above $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$.

Let us compute $\dim \mathcal{B}_{\sigma'}$. It is enough to compute it in one of the cases since one case can be obtained from the other by mirroring P_σ around the point $\frac{1+n}{2}$.

For example, let us consider case $\sigma' = \sigma_{\curvearrowright(i,j)}$. Let $\sigma' = \sigma_{i \rightarrow m}$. Recall notation from the proof of Theorem 3.1. Note that

- $f'_{(m,j)}(P_{\sigma'}) = f'_{(i,j)}(P_{\sigma}) + 1$ and since $\mathbf{over}_{(i,j)}(P_{\sigma}) \subset \mathbf{over}_m(P_{\sigma})$ one has $c'_{(m,j)}(P_{\sigma'}) = c'_{(i,j)}(P_{\sigma})$.
- Again, since $\mathbf{over}_{(i,j)}(P_{\sigma}) \subset \mathbf{over}_m(P_{\sigma})$ one has $\mathbf{over}_{(m,j)}(P_{\sigma'}) = \mathbf{over}_{(i,j)}(P_{\sigma})$ so that for any $(i', j') \in \mathbf{over}_{(i,j)}(P_{\sigma})$ one has $f'_{(i',j')}(P_{\sigma'}) = f'_{(i',j')}(P_{\sigma})$ and $c'_{(i',j')}(P_{\sigma'}) = c'_{(i',j')}(P_{\sigma})$.
- Exactly in the same way for $(i', j') \in P_{\sigma}$ such that $(i', j') \notin \mathbf{over}_m(P_{\sigma})$ one has $f'_{(i',j')}(P_{\sigma'}) = f'_{(i',j')}(P_{\sigma})$ and $c'_{(i',j')}(P_{\sigma'}) = c'_{(i',j')}(P_{\sigma})$.
- If $(i', j') \in (\mathbf{over}_m(P_{\sigma}) \setminus \mathbf{over}_{(i,j)}(P_{\sigma})) \cap c'_{(i,j)}(P_{\sigma})$ (that is $i' < m$ and $i < j' < j$) then again $f'_{(i',j')}(P_{\sigma'}) = f'_{(i',j')}(P_{\sigma})$ and $c'_{(i',j')}(P_{\sigma'}) = c'_{(i',j')}(P_{\sigma})$.
- Finally, if $(i', j') \in (\mathbf{over}_m(P_{\sigma}) \setminus \mathbf{over}_{(i,j)}(P_{\sigma})) \cap l_{(i,j)}(P_{\sigma})$ (that is $i' < m$ and $m < j' < i$) then $f'_{(i',j')}(P_{\sigma'}) = f'_{(i',j')}(P_{\sigma}) - 1$ and $c'_{(i',j')}(P_{\sigma'}) = c'_{(i',j')}(P_{\sigma}) + 1$.

Set $t := |(\mathbf{over}_m(P_{\sigma}) \setminus \mathbf{over}_{(i,j)}(P_{\sigma})) \cap l_{(i,j)}(P_{\sigma})|$. Summarizing the calculations above we get

$$\begin{aligned} f(P_{\sigma'}) &= \sum_{s=1}^k f'_{(i_s, j_s)}(P_{\sigma'}) = f(P_{\sigma}) + 1 - t; \\ c(P_{\sigma'}) &= \sum_{s=1}^k c'_{(i_s, j_s)}(P_{\sigma'}) = c(P_{\sigma}) + t; \end{aligned}$$

so that $\dim \mathcal{B}_{\sigma'} = k(n - k) - f(P_{\sigma'}) - c(P_{\sigma'}) = \dim \mathcal{B}_{\sigma} - 1$. $\square$

3.5. Let us again consider $\sigma = (i_1, j_1) \dots (i_k, j_k)$. Take $i, j \in \langle \sigma \rangle$ such that $i < j$. Assume that they belong to different pairs, that is $i \in (i_s, j_s)$ and $j \in (i_t, j_t)$ where $s \neq t$. Put $\sigma_{i \rightleftharpoons j}$ to be an involution obtained by interchanging places of i and j in the pairs. Assume that $(i_s, j_s) = ((i, p))$ and $(i_t, j_t) = ((j, q))$ then $\sigma_{i \rightleftharpoons j} = ((i, q))((j, p)) \dots$ where by dots we denote all the pairs of σ but $(i_s, j_s), (i_t, j_t)$. Note that we cannot say anything about ordering $((i, q))$ and $((j, p))$ so we write them in double round brackets.

The link pattern $P_{\sigma'}$ described below is obtained from P_{σ} by crossing some arc with an arc to the left of it.

Let $(i, j) \in \sigma$. Translating [11, 3.13] we put

$$L_{(i,j)}(\sigma) := \{(i_s, j_s) \in \sigma \mid j_s < i \text{ and } \{p\}_{p=j_s+1}^{i-1} \subset \langle \pi_{i_s, j_s}(P_{\sigma}) \rangle\}$$

In other words $(i_s, j_s) \in L_{(i,j)}(\sigma)$ if $(i_s, j_s) \in l_{(i,j)}(P_{\sigma})$ and there are no fixed points between j_s and i in $\pi_{i_s, j_s}(P_{\sigma})$. Note that $L_{(i,j)}(\sigma)$ may contain a few elements. Put


$$S_{(i,j) \rightarrow}(\sigma) := \begin{cases} \{\sigma_{j_s \rightleftharpoons i} \mid (i_s, j_s) \in L_{(i,j)}(\sigma)\} & \text{if } L_{(i,j)}(\sigma) \neq \emptyset \\ \emptyset & \text{otherwise} \end{cases}$$

Note that for $(i_s, j_s) \in L_{(i,j)}(\sigma)$ one has


$$(R_{\sigma_{j_s \rightleftharpoons i}})_{s,t} = \begin{cases} (R_{\sigma})_{s,t} - 1 & \text{if } i \leq s < j_s \text{ and } t \geq j \text{ or } j_s \leq t < i \text{ and } s \leq i_s; \\ (R_{\sigma})_{s,t} & \text{otherwise.} \end{cases}$$

Thus, for any $(i_s, j_s) \in L_{(i,j)}(\sigma)$ one has $\sigma_{j_s \rightleftharpoons i} \prec \sigma$.


For example, let $\sigma = (1, 5)(2, 4)(3, 6)(7, 9)(10, 11) \in \mathbf{S}_{11}^2$. Its link pattern is



One has $L_{(1,5)}(\sigma) = L_{(2,4)}(\sigma) = L_{(3,6)}(\sigma) = \emptyset$, $L_{(7,9)}(\sigma) = \{(1,5), (3,6)\}$, so that



and $L_{(10,11)}(\sigma) = \{(7,9)\}$ so that



Again, as an easy corollary of Theorem 3.1 we get

Proposition. *If $L_{(i,j)}(\sigma) \neq \emptyset$ then $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$ for any $\sigma' \in S_{(i,j) \rightarrow}(\sigma)$*

Proof. By the note above $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$. Also, by the definition of $L_{(i,j)}(\sigma)$ $P_{\sigma'}$ differs from P_σ by one additional cross obtained from changing pairs $(i_s, j_s)(i, j)$ to $(i_s, i)(j_s, j)$ as we show below.

Indeed, since there are no fixed points between j_s and i in $\pi_{i_s, j}(P_\sigma)$ we get that in particular there are no fixed points between j_s and i in P_σ so that $f(P_\sigma) = f(P_{\sigma'})$. Also, the absence of fixed points between j_s and i in $\pi_{i_s, j}(P_\sigma)$ means that $\mathbf{over}_{(i_s, j_s)}(P_\sigma) \cap l_{(i, j)}(P_\sigma) = \emptyset$ and $\mathbf{over}_{(i, j)}(P_\sigma) \cap r_{(i_s, j_s)}(P_\sigma) = \emptyset$ so that $c(P_{\sigma'}) = c(P_\sigma) + 1$. Thus, by Theorem 3.1 $\dim \mathcal{B}_{\sigma'} = \dim \mathcal{B}_\sigma - 1$. $\square$

3.6. The last type of elements in $\mathcal{D}(\sigma)$ is obtained by crossing two concentric arcs. It is defined as follows. For $(i, j) \in \sigma$ recall $\mathbf{over}_{(i, j)}(P_\sigma)$ and $\mathbf{under}_{(i, j)}(P_\sigma)$ from 3.1 and put

$$Ov_{(i, j)}(\sigma) := \{(i_s, j_s) \in \mathbf{over}_{(i, j)}(P_\sigma) \mid \mathbf{under}_{(i_s, j_s)}(P_\sigma) = \{(i, j)\} \cup \mathbf{under}_{(i, j)}(P_\sigma)\}$$

In other words $(i_s, j_s) \in Ov_{(i,j)}(\sigma)$ if $(i_s, j_s) \in \mathbf{over}_{(i,j)}(P_\sigma)$ and there are no arcs between (i_s, j_s) and (i, j) in P_σ . Note that $Ov_{(i,j)}(\sigma)$ may contain a few elements. Put


$$S_{(i,j)1\downarrow}(\sigma) := \begin{cases} \{\sigma_{i_s \leftrightsquigarrow i} \mid (i_s, j_s) \in Ov_{(i,j)}(\sigma)\} & \text{if } Ov_{(i,j)}(\sigma) \neq \emptyset; \\ \emptyset & \text{otherwise;} \end{cases}$$

Note that $\sigma_{i_s \leftrightsquigarrow i_t} = \sigma_{j_s \leftrightsquigarrow j_t}$. Note also that for $(i_p, j_p) \in Ov_{(i,j)}(\sigma)$ one has


$$(R_{\sigma_{i_p \leftrightsquigarrow i}})_{s,t} = \begin{cases} (R_\sigma)_{s,t} - 1 & \text{if } i_p < s \leq i \text{ and } j \leq t < j_s; \\ (R_\sigma)_{s,t} & \text{otherwise.} \end{cases}$$

Thus, for any $(i_p, j_p) \in Ov_{(i,j)}(\sigma)$ one has $\sigma_{i_p \leftrightsquigarrow i} \prec \sigma$.


For example, let $\sigma = (1, 11)(2, 6)(3, 9)(4, 5) \in \mathbf{S}_{11}^2$. Its link pattern is




One has $Ov_{(1,11)}(\sigma) = \emptyset$, $Ov_{(2,6)}(\sigma) = \{(1, 11)\}$ so that



and $Ov_{(3,9)}(\sigma) = \{(1, 11)\}$ so that



Finally $Ov_{(4,5)}(\sigma) = \{(2, 6), (3, 9)\}$ so that



And again, exactly as in 3.5 we get

Proposition. *If $Over_{(i,j)}(\sigma) \neq \emptyset$ then $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$ for any $\sigma' \in S_{(i,j)1\downarrow}(\sigma)$*

Proof. Exactly as in 3.5 we get that $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ and exactly as in 3.5 we get that $f(P_{\sigma'}) = f(P_\sigma)$ and $c(P_{\sigma'}) = c(P_\sigma) + 1$ which together provide us the result. $\square$

3.7. By Proposition 3.15 from [11] and Propositions 3.4, 3.5, 3.6 we get

Theorem. *Let $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2(k)$ then*

- (i) $\mathcal{D}(\sigma) = \coprod_{s=1}^k \sigma_{(i_s, j_s) \curvearrowright} \sqcup \coprod_{s=1}^k \sigma_{\curvearrowleft (i_s, j_s)} \sqcup \coprod_{s=1}^k S_{(i_s, j_s) \curvearrowright}(\sigma) \sqcup \coprod_{s=1}^k S_{(i_s, j_s) \downarrow}(\sigma)$
- (ii) *Let $\sigma' \in \mathbf{S}_n^2(k)$ be such that $\sigma' \prec \sigma$. Then $\sigma' \in \mathcal{D}(\sigma)$ iff $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$.*

3.8. Now we are ready to finish the description of $\mathcal{C}(\sigma)$. Note first that

$$\mathcal{D}(\sigma) \subset \mathcal{C}(\sigma) \subset \mathcal{D}(\sigma) \cup \mathcal{N}(\sigma)$$

So we have to find $\mathcal{N}(\sigma) \cap \mathcal{C}(\sigma)$.

Consider $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$. Let $E_{\max}(\sigma)$ be the subset of arcs in $E(\sigma)$ without fixed points outside of them. In that case all the fixed points of P_σ are under arc (i, j) so that $(i, j) \in E_{\max}(\sigma)$ iff $f'_{(i,j)}(P_\sigma) = n - 2k$. We can use this fact to define $E_{\max}(\sigma)$ formally:

$$E_{\max}(\sigma) := \{(i, j) \in E(\sigma) \mid f'_{(i,j)}(P_\sigma) = n - 2k\}$$

Lemma. *Consider $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$ and let $(i, j) \in E(\sigma)$.*

- (i) *If $(i, j) \notin E_{\max}(\sigma)$ then $\sigma_{(i,j)}^- \notin \mathcal{C}(\sigma)$.*
- (ii) *If $(i, j) \in E_{\max}(\sigma)$ then $\dim \mathcal{B}_{\sigma_{(i,j)}^-} = \dim \mathcal{B}_\sigma - 1$.*

Proof. To prove (i) note that for any $(i, j) \in E(\sigma)$ one has $\text{over}_{(i,j)}(P_\sigma) = \emptyset$. So the conditions of 3.4 for $(i, j) \in E(\sigma)$ are satisfied iff $(i, j) \notin E_{\max}(\sigma)$ and in this case $S = \{\sigma_{\curvearrowleft (i,j)}, \sigma_{(i,j) \curvearrowright}\} \neq \emptyset$. Let $\sigma' \in S$ and let $P_{\sigma'}$ be obtained from P_σ by

replacing (i, j) to (i', j') . Then obviously $\sigma_{(i,j)}^- = (\sigma')_{(i',j')}^-$ so that $\sigma_{(i,j)}^- \prec \sigma' \prec \sigma$ and $\sigma_{(i,j)}^- \notin \mathcal{C}(\sigma)$.

To compute (ii) we use notation from the proof of theorem 3.1. Consider an arc $(i_s, j_s) \in P_\sigma$.

- If (i_s, j_s) does not intersect (i, j) then

$$f'_{(i_s, j_s)}(P_\sigma) = f'_{(i_s, j_s)}(P_{\sigma_{(i,j)}^-}) \quad \text{and} \quad |c_{(i_s, j_s)}^l(P_\sigma)| = |c_{(i_s, j_s)}^l(P_{\sigma_{(i,j)}^-})|.$$

- If (i_s, j_s) intersects (i, j) on the left i is a new fixed point under (i_s, j_s) in $P_{\sigma_{(i,j)}^-}$ comparing to P_σ and the crossings from the left of (i_s, j_s) are the same in both patterns, so that

$$f'_{(i_s, j_s)}(P_\sigma) = f'_{(i_s, j_s)}(P_{\sigma_{(i,j)}^-}) - 1 \quad \text{and} \quad |c_{(i_s, j_s)}^l(P_\sigma)| = |c_{(i_s, j_s)}^l(P_{\sigma_{(i,j)}^-})|$$

- If (i_s, j_s) intersects (i, j) on the right then j is a new fixed point under (i_s, j_s) in $P_{\sigma_{(i,j)}^-}$ comparing to P_σ and there is one less crossing from the left of (i_s, j_s) in $P_{\sigma_{(i,j)}^-}$ comparing to P_σ , so that

$$f'_{(i_s, j_s)}(P_\sigma) = f'_{(i_s, j_s)}(P_{\sigma_{(i,j)}^-}) - 1 \quad \text{and} \quad |c_{(i_s, j_s)}^l(P_\sigma)| = |c_{(i_s, j_s)}^l(P_{\sigma_{(i,j)}^-})| + 1$$

- Finally, since there are no fixed points outside of (i, j) in P_σ one has $f'_{(i,j)}(P_\sigma) = n - 2k$

Summarizing we get

$$\begin{aligned} f(P_\sigma) &= \sum_{s=1}^k f'_{(i_s, j_s)}(P_\sigma) = \sum_{(i_s, j_s) \neq (i, j)} f'_{(i_s, j_s)}(P_\sigma) + f'_{(i, j)}(P_\sigma) = \\ &= \sum_{(i_s, j_s) \neq (i, j)} f'_{(i_s, j_s)}(P_{\sigma_{(i,j)}^-}) - |c_{(i, j)}^l(P_\sigma)| - |c_{(i, j)}^r(P_\sigma)| + n - 2k = \\ &= f(P_{\sigma_{(i,j)}^-}) - |c_{(i, j)}^l(P_\sigma)| - |c_{(i, j)}^r(P_\sigma)| + n - 2k; \\ c(P_\sigma) &= \sum_{s=1}^k |c_{(i_s, j_s)}^l(P_\sigma)| = \sum_{(i_s, j_s) \neq (i, j)} |c_{(i_s, j_s)}^l(P_\sigma)| + |c_{(i, j)}^l(P_\sigma)| = \\ &= \left(\sum_{\substack{(i_s, j_s) \neq (i, j) \\ (i_s, j_s) \notin c_{(i, j)}^r(P_\sigma)}} |c_{(i_s, j_s)}^l(P_{\sigma_{(i,j)}^-})| \right) + \left(\sum_{(i_s, j_s) \in c_{(i, j)}^r(P_\sigma)} |c_{(i_s, j_s)}^l(P_{\sigma_{(i,j)}^-})| + |c_{(i, j)}^r(P_\sigma)| \right) + |c_{(i, j)}^l(P_\sigma)| = \\ &= c(P_{\sigma_{(i,j)}^-}) + |c_{(i, j)}^r(P_\sigma)| + |c_{(i, j)}^l(P_\sigma)| \end{aligned}$$

Thus,

$$\begin{aligned}
\dim \mathcal{B}_\sigma &= k(n-k) - f(P_\sigma) - c(P_\sigma) = \\
&= k(n-k) - \left(f(P_{\sigma_{(i,j)}^-}) - |c_{(i,j)}^l(P_\sigma)| - |c_{(i,j)}^r(P_\sigma)| + n - 2k \right) - \\
&\quad \left(c(P_{\sigma_{(i,j)}^-}) + |c_{(i,j)}^r(P_\sigma)| + |c_{(i,j)}^l(P_\sigma)| \right) = \\
&= k(n-k) - (n-2k) - f(P_{\sigma_{(i,j)}^-}) - c(P_{\sigma_{(i,j)}^-}) = \\
&= (k-1)(n-k+1) + 1 - f(P_{\sigma_{(i,j)}^-}) - c(P_{\sigma_{(i,j)}^-}) = \dim \mathcal{B}_{\sigma_{(i,j)}^-} + 1.
\end{aligned}$$

□

3.9. As a straightforward corollary of theorem 3.7 and lemma 3.8, we get

Theorem. Let $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$. One has

- (i) $\mathcal{C}(\sigma) = \mathcal{D}(\sigma) \sqcup \{\sigma_{(i,j)}^- \mid (i,j) \in E_{\max}(\sigma)\}$.
- (ii) $\sigma' \in \mathcal{C}(\sigma)$ iff $\text{codim}_{\overline{\mathcal{B}}_\sigma} \mathcal{B}_{\sigma'} = 1$.

3.10. In section 5 we will discuss intersections of **B** orbit closures in $\mathcal{X}_n$. As a preparation we would like to finish this section with a few preliminary general definition and results.

Given $\emptyset \neq S \subset \mathbf{S}_n^2$. We define the components of $U = \bigcap_{\sigma \in S} \overline{\mathcal{B}}_\sigma$ in a standard way by setting $\mathcal{B}_{\tau_1}, \dots, \mathcal{B}_{\tau_r}$ to be the components of U if $U = \bigcup_{i=1}^r \overline{\mathcal{B}}_{\tau_i}$ and $\tau_i \preceq \tau_j$ implies $i = j$ for any $i, j : 1 \leq i, j \leq r$. Let us call $\tau_1, \dots, \tau_r$ component involutions of U .

Given $\sigma \in \mathbf{S}_n^2(k)$ where $k \leq \frac{1}{2}n - 1$. For $m : k < m \leq \frac{1}{2}n$ put $F_m(\sigma) := \{\tau \in \mathbf{S}_n^2(m) \mid \tau \succ \sigma\}$. Let $\{p_i\}_{i=1}^{n-2k}$ be fixed points of σ . Note that $n - 2k \geq 2(m - k)$ so that $\sigma' = \sigma(p_1, p_2) \dots (p_{2(m-k)-1}, p_{2(m-k)}) \in \mathbf{S}_n^2(m)$ and $\sigma' \succ \sigma$. Thus, $F_m(\sigma) \neq \emptyset$

As an easy corollary of subsection 2.6 we get

Proposition. Let $k \leq \frac{1}{2}n - 1$. For any $\sigma \in \mathbf{S}_n^2(k)$ and any $m : k < m \leq \frac{1}{2}n$ there exists the unique $\overline{\sigma}_m \in F_m(\sigma)$ such that

$$\bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau = \overline{\mathcal{B}}_{\overline{\sigma}_m}.$$

In particular, let $\emptyset \neq S \subset \mathbf{S}_n^2(k)$ and let $U = \bigcap_{\sigma \in S} \overline{\mathcal{B}}_\sigma$. Then all the component involutions of U are of length k .

Proof. By (*) of 2.6 $U' = \bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau \cap \mathcal{O}_{(n-m,m)^*} \neq \emptyset$. Let $\tau_1, \dots, \tau_r$ be the component involutions of U' . Then by irreducibility of $\mathcal{B}_\sigma$ there exists $\tau_i = \overline{\sigma}_m$ such that $\mathcal{B}_\sigma \subset \overline{\mathcal{B}}_{\overline{\sigma}_m}$ but then $\overline{\sigma}_m \in F_m(\sigma)$ and

$$U' = \bigcup_{j=1}^r (\overline{\mathcal{B}}_{\tau_j} \cap \overline{\mathcal{B}}_{\overline{\sigma}_m} \cap \mathcal{O}_{(n-m,m)^*}) = \overline{\mathcal{B}}_{\overline{\sigma}_m} \cap \mathcal{O}_{(n-m,m)^*}.$$

To complete the proof we have to show that $\bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau = \overline{\mathcal{B}}_{\overline{\sigma}_m}$ (in other words we have to show that $\overline{\bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau \cap \mathcal{O}_{(n-m,m)^*}} = \bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau \cap \overline{\mathcal{O}_{(n-m,m)^*}} = \bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau$, where the second equality is obvious). To do this we must show that for any $j < m$ for any $\mathcal{B}_{\sigma'} \subset \bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau \cap \mathcal{O}_{(n-j,j)^*}$ one has $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_{\overline{\sigma}_m}$. Let us consider $F_m(\sigma')$. Since $\mathcal{B}_{\sigma'} \subset \bigcap_{\tau \in F_m(\sigma)} \overline{\mathcal{B}}_\tau$ we get $F_m(\sigma) \subset F_m(\sigma')$. By the previous there exists the unique minimal element $\overline{\sigma}'_m$ of $F_m(\sigma')$ so that $\overline{\sigma}'_m \preceq \overline{\sigma}_m$ which provides $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_{\overline{\sigma}_m}$ and completes the proof.

Let us consider U from the proposition. It is enough to show that for any σ such that $\mathcal{B}_\sigma \subset U$ and $l(\sigma) < k$ there exists $\sigma' \succ \sigma$ such that $\mathcal{B}_{\sigma'} \subset U$ and $l(\sigma') = k$. Consider $F_k(\sigma)$. By the definition of U we get $S \subset F_k(\sigma)$ so that $\overline{\sigma}_k$ satisfies $\mathcal{B}_{\overline{\sigma}_k} \subset U$. $\square$

Note that in general U defined in the proposition can be reducible.

3.11. Given $\sigma' \in F_p(\sigma)$, to show that $\sigma' = \overline{\sigma}_p$ it is enough to check that


$$\dim \mathcal{B}_{\sigma''} \geq \dim \mathcal{B}_{\sigma'} \quad \text{for any } \sigma'' \in F_p(\sigma). \quad (*)$$

It is easy to write explicitly $\overline{\sigma}_{k+1}$ and to compute the dimension of $\mathcal{B}_{\overline{\sigma}_{k+1}}$. For $\sigma \in \mathbf{S}_n^2(k)$ let $i_\sigma := \min\{\{s\}_{s=1}^n \setminus \langle \sigma \rangle\}$ and $j_\sigma := \max\{\{s\}_{s=1}^n \setminus \langle \sigma \rangle\}$. If $E_{\max}(\sigma) \neq \emptyset$ then $(i, j) \in \text{over}_{i_\sigma}(P_\sigma), \text{over}_{j_\sigma}(P_\sigma)$ for any $(i, j) \in E_{\max}$. Let $E_{\max}(\sigma) = \{(i_r, j_r)\}_{r=1}^s$ where the arcs are ordered in the increasing order according to the left end points. Note that $i_1 < i_2 < \dots < i_s < i_\sigma < j_\sigma < j_1 < \dots < j_s$. Put $j_0 := j_\sigma$ and $i_{s+1} := i_\sigma$ and let σ' be obtained from σ by deleting all the cycles in $E_{\max}(\sigma)$. Put


$$\overline{\sigma}_{k+1} = \begin{cases} \sigma(i_\sigma, j_\sigma) & \text{if } E_{\max}(\sigma) = \emptyset \\ \sigma'(i_1, j_0)(i_2, j_1) \dots (i_s, j_{s-1})(i_{s+1}, j_s) & \text{otherwise} \end{cases}$$

Let us show some examples to make the picture clear:

$E_{\max}(\sigma) = \emptyset$:



$$E_{\max}(\sigma) = \{(2, 10)\} :$$



Note that

$$(R_{\bar{\sigma}_{k+1}})_{i,j} = \begin{cases} (R_{\sigma})_{i,j} + 1 & \text{if } i \leq i_r \text{ and } j \geq j_{r-1} \text{ for } 1 \leq r \leq s \\ (R_{\sigma})_{i,j} & \text{otherwise} \end{cases}$$

So that $\bar{\sigma}_{k+1} \succ \sigma$. We must show that $\bar{\sigma}_{k+1}$ defined above is indeed the minimal element of $F_{k+1}(\sigma)$.

Proposition. *Let $k \leq \frac{1}{2}n - 1$. Then for every $\sigma' \in F_p(\sigma)$ where $p \geq k + 1$ one has $\sigma' \succeq \bar{\sigma}_{k+1}$. One also has $\text{codim}_{\bar{\mathcal{B}}_{\bar{\sigma}_{k+1}}} \mathcal{B}_{\sigma} = |E_{\max}(\sigma)| + 1$.*

Proof. Let us show this by induction on the $|E_{\max}(\sigma)|$. Put $A(\sigma) := \{\sigma' \mid \sigma \in \mathcal{D}(\sigma')\}$ and $B(\sigma) := \{\sigma' \mid \sigma \in \mathcal{C}(\sigma') \text{ and } \ell(P_{\sigma'}) = \ell(P_{\sigma}) + 1\}$. Note that for any $\sigma' \in A(\sigma)$ one has

$$|E_{\max}(\sigma)| - 1 \leq |E_{\max}(\sigma')| \leq |E_{\max}(\sigma)|. \quad (**)$$

Note also that by definition

$$B(\sigma) = \begin{cases} \sigma(i_{\sigma}, j_{\sigma}) & \text{if } E_{\max}(\sigma) = \emptyset \\ \emptyset & \text{otherwise} \end{cases} \quad (**_*)$$

If $E_{\max}(\sigma) = \emptyset$ then $B(\sigma) = \{\sigma(i_{\sigma}, j_{\sigma})\}$ so that $\text{codim}_{\bar{\mathcal{B}}_{\sigma(i_{\sigma}, j_{\sigma})}} \mathcal{B}_{\sigma} = 1$. For any $\sigma' \in F_{k+1}(\sigma)$ one has $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma} \geq 1$, thus, by (*) and Proposition 3.11 we get the result.

Now assume this is true for $\sigma \in \mathbf{S}_n^2(k)$ such that $|E_{\max}(\sigma)| = s$ and show this is true for σ such that $|E_{\max}(\sigma)| = s + 1$. Let σ' be obtained from σ by changing (i_1, j_1) to (i_1, j_{σ}) . Note that $\sigma = \sigma'_{(i_1, j_{\sigma}) \curvearrowright}$ so that $\sigma' \in A(\sigma)$ and that $|E_{\max}(\sigma')| = s$. Note also that $\bar{\sigma}_{k+1} = \bar{\sigma}'_{k+1}$, thus, by induction one has $\text{codim}_{\bar{\mathcal{B}}_{\bar{\sigma}_{k+1}}} \mathcal{B}_{\sigma} = \text{codim}_{\bar{\mathcal{B}}_{\bar{\sigma}_{k+1}}} \mathcal{B}_{\sigma'} + \text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma} = s + 1$.

To show that $\bar{\sigma}_{k+1}$ is the minimal element of $F_{k+1}(\sigma)$ it is enough to show by (*) that $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma} \geq s + 1$ for any $\sigma' \in F_{k+1}(\sigma)$. Assume this is not true. Let $\sigma \in \mathbf{S}_n^2(k)$ be maximal such that $|E_{\max}(\sigma)| = s + 1$ and there exists $\sigma' \in F_{k+1}(\sigma)$ such that $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma} \leq s$. By (**_{*}) $B(\sigma) = \emptyset$ so that by theorem 3.9 there exists $\sigma'' \in A(\sigma)$ such that $\sigma' \succ \sigma'' \succ \sigma$. Note that $\sigma' \in F_{k+1}(\sigma'')$. By (**) $s \leq |E_{\max}(\sigma'')| \leq s + 1$. If $|E_{\max}(\sigma'')| = s + 1$ we get that $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma''} \leq s - 1$ which contradicts maximality of σ . If $|E_{\max}(\sigma'')| = s$ then by induction hypothesis $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma''} \geq s$ so that $\text{codim}_{\bar{\mathcal{B}}_{\sigma'}} \mathcal{B}_{\sigma} \geq s + 1$ which contradicts the assumption. $\square$

4. COMBINATORIAL DESCRIPTION OF THE CLOSURE OF AN ORBITAL VARIETY OF NILPOTENT ORDER 2

4.1. In this section we apply the results obtained above to the closures of $\mathbf{B}$ orbits of dimension $k(n-k)$ in $\mathfrak{B}_{(n-k,k)}$, that is of the maximal possible dimension. Such orbit is dense in an orbital variety associated to $\mathcal{O}_{(n-k,k)^*}$ and every orbital variety associated to $\mathcal{O}_{(n-k,k)^*}$ admits such $\mathbf{B}$ orbit as we explained in 2.2.

As it was shown in [11, 4.5] the closure of an orbital variety of nilpotent order 2 is a union of orbital varieties. In [11, 4.3] the combinatorial description of such orbital variety in terms of Young tableaux is given. We would like to use link patterns to simplify this description.

By Proposition 2.2 $\mathcal{B}_T$ is the dense $\mathbf{B}$ -orbit in $\mathcal{V}_T$ for $T \in \mathbf{Tab}_{(n-k,k)^*}$. By definition of an orbital variety $\overline{\mathcal{V}}_T = \mathcal{V}_T \sqcup \bigcup_{\sigma' \in \mathcal{N}(\sigma_T)} \overline{\mathcal{B}}_{\sigma'}$. Moreover, since an element of $\mathcal{N}(\sigma_T)$ is

obtained from σ_T by erasing an external arc it is a link pattern without fixed points under an arc and without arc crossings, thus, $\mathcal{B}_{\sigma'}$ for $\sigma' \in \mathcal{N}_{\sigma_T}$ is $\sigma' = \sigma_{T'}$ for some $T' \in \mathbf{Tab}_{(n-k+1,k-1)^*}$.

How to define $\{T' \in \mathbf{Tab}_{(n-k+1,k-1)^*} \mid \sigma_{T'} \in \mathcal{N}(\sigma_T)\}$? The answer given in [11, 4.3] is too complex to be satisfactory. Let us try to understand the picture using link patterns.

Given $\sigma_T = (i_1, j_1) \dots (i_k, j_k)$ ordered in increasing order according to the second entry of the cycle (that is $j_1 < \dots < j_k$). Then $T_2 = \begin{pmatrix} j_1 \\ \vdots \\ j_k \end{pmatrix}$ (cf. 2.2). Moreover, set $\{j_1, \dots, j_k\}$ defines σ_T completely. In turn $\sigma_{T'}$ is obtained from σ_T by erasing

some external arc, that is $T'_2 = \begin{pmatrix} \vdots \\ j_{l-1} \\ j_{l+1} \\ \vdots \end{pmatrix}$ for some $l : 1 \leq l \leq k$. To determine

$T' : \sigma_{T'} \in \mathcal{N}_{\sigma_T}$ we must find $E(\sigma_T)$ – the subset of external arcs.

Note that since arcs of P_{σ_T} do not intersect, for any $t > s$ (that is such that $j_t > j_s$) one has $i_t \notin [i_s, j_s]$ that is either $i_t < i_s$ or $i_t > j_s$. Thus, arc $(i_s, j_s) \in E(\sigma_T)$ if for any $(i_t, j_t) \in \sigma$ such that $t > s$ one has $i_t > j_s$. Hence, for any $t > s$ there must be at least $2(t-s) - 1$ points between j_s and j_t . In other words the necessary condition is $j_t - j_s \geq 2(t-s)$. This is also a sufficient condition. Let us formulate this statement as a lemma

Lemma. Let $T \in \mathbf{Tab}_{(n-k,k)^*}$ be such that $T_2 = \begin{pmatrix} j_1 \\ \vdots \\ j_k \end{pmatrix}$. Let $\sigma_T = (i_1, j_1) \dots (i_k, j_k)$.

Then $E(\sigma_T) = \{(i_l, j_l) \mid l = k \text{ or } \forall s : l < s \leq k \Rightarrow j_s - j_l \geq 2(s-l)\}$.

Proof. First of all let us show that if $(i_l, j_l) \in E(\sigma_T)$ then for any $s > l$ one must have $j_s - j_l \geq 2(s - l)$. Indeed, for any $t : l < t \leq s$ one has (as we already mentioned) $i_t > j_l$ and $j_t \leq j_s$ so that $[j_l + 1, j_s]$ must contain at least $2(s - l)$ points, therefore $j_s - j_l \geq 2(s - l)$.

Exactly in the same way we show that if $j_s - j_l \geq 2(s - l)$ for any $s > l$ then $(i_l, j_l) \in E(\sigma_T)$. Indeed, for $s = l + 1$ one has $j_{l+1} \geq j_l + 2$ so that $j_{l+1} - 1 \in \langle T_1 \rangle$ and $i_{l+1} = j_{l+1} - 1 > j_l$. Assume that we already know arcs $(i_{l+1}, j_{l+1}), \dots, (i_{s-1}, j_{s-1})$ and $i_{l+1}, \dots, i_{s-1} \in [j_l + 1, j_{s-1}]$. By the construction $i_s = \max\{r\}_{r=1}^n \setminus \{i_p, j_p\}_{p=1}^{s-1}$. And since there are at least $2(s - l)$ points in $[j_l + 1, j_s]$ and only $2(s - 1 - l) + 1$ are occupied we get that $i_s \in [j_l + 1, j_s]$. $\square$

4.2. Consider $T \in \mathbf{Tab}_{(n-k,k)^*}$ and let $T_2 = \begin{pmatrix} j_1 \\ \vdots \\ j_k \end{pmatrix}$. For $b \in \langle T_2 \rangle$ put $T\langle b \rangle$ to be

the tableau obtained from T by moving b from T_2 to an appropriate place in T_1 .

Let us denote by $\mathcal{N}(T) := \{S \in \mathbf{Tab}_{(n-k+1,k-1)^*} \mid \mathcal{V}_S \subset \mathcal{V}_T\}$. By dimension reasoning it is obvious that $\mathcal{V}_S : S \in \mathcal{N}(T)$ are orbital varieties of maximal dimension in $\overline{\mathcal{V}}_T \setminus \mathcal{V}_T$. Moreover the fact that $\mathcal{V}_T$ admits a dense $\mathbf{B}$ orbit and Proposition 3.3 provide that these are all maximal orbital varieties in $\overline{\mathcal{V}}_T \setminus \mathcal{V}_T$. In other words $\overline{\mathcal{V}}_T = \mathcal{V}_T \sqcup \bigcup_{S \in \mathcal{N}(T)} \overline{\mathcal{V}}_S$.

As a straightforward corollary of Lemma 4.1 we get

Theorem. Let $T \in \mathbf{Tab}_{(n-k,k)^*}$ and $T_2 = \begin{pmatrix} j_1 \\ \vdots \\ j_k \end{pmatrix}$. Then

$$\mathcal{N}(T) = \{T\langle j_i \rangle \mid i = k \text{ or } \forall s : i < s \leq k \Rightarrow j_s - j_i \geq 2(s - i)\}.$$

This theorem simplifies Theorem 4.3 from [11].

5. MEANDERS AND INTERSECTIONS OF THE COMPONENTS OF A SPRINGER FIBER OF NILPOTENT ORDER 2


5.1. In this section we apply our results to the theory of intersections of the components of Springer fiber $\mathcal{F}_x$ where $x \in \mathcal{O}_{(n-k,k)^*}$. For $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ let $\mathcal{F}_T, \mathcal{F}_S$ be the corresponding components of $\mathcal{F}_x$. As we explained in 1.9 the number of components of $\mathcal{F}_T \cap \mathcal{F}_S$ and their codimensions equal to the number of components of $\mathcal{V}_T \cap \mathcal{V}_S$ and their codimensions. So we formulate our results in terms of the components of a Springer fiber, however the computations are made in $\mathbf{B}$ orbits of nilpotent order 2. First note that for $T, S \in \mathbf{Tab}_{(n-k,k)^*}$ one has

$$\mathcal{V}_T \cap \mathcal{V}_S = \overline{\mathcal{B}}_T \cap \overline{\mathcal{B}}_S \cap \mathcal{O}_{(n-k,k)^*} = \coprod_{\substack{\sigma \in \mathbf{S}_n^2(k) \\ \sigma \preceq \sigma_T, \sigma_S}} \mathcal{B}_\sigma.$$

We will try to understand these intersections in terms of meanders. We draw one link pattern upwards and one link pattern downwards. If both link patterns of a meander are without intersections we get a classical meander, otherwise we get a generalized meander.

As we have mentioned in 1.9 a generalized meander $M_{\sigma,\sigma'}$ corresponding to $\sigma, \sigma' \in \mathbf{S}_n^2(k)$ provides us an easy way to compute $R_{\sigma,\sigma'}$. However for a generalized meander the picture seems to be too complex to understand it without going to matrix $R_{\sigma,\sigma'}$.

As it is shown in [14, 3.8] even an intersection of codimension 1 for (closures of) generic $\mathbf{B}$ orbits of the same dimension can be reducible. Moreover, it can be not of pure dimension as it is shown by the example below. We also use this example to construct the corresponding generalized meander. Consider $\sigma = (1, 3)(4, 5)$, $\sigma' = (2, 3)(4, 6) \in \mathbf{S}_6^2(2)$. Note that $\dim \mathcal{B}_{(1,3)(4,5)} = \dim \mathcal{B}_{(2,3)(4,6)} = 2 \cdot 4 - 1 = 7$. The generalized meander is



and

$$R_{(1,3)(4,5), (2,3)(4,6)} = \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

One can easily check using Proposition 2.3 that $R_{(1,3)(4,5), (2,3)(4,6)} \notin \mathbf{R}_6^2$ so that the intersection is reducible by Theorem 2.6. Computing $R_{(1,3)(4,6)}$ and $R_{(1,6)(2,5)}$ we get $\overline{\mathcal{B}}_{(1,3)(4,5)} \cap \overline{\mathcal{B}}_{(2,3)(4,6)} = \overline{\mathcal{B}}_{(1,3)(4,6)} \cup \overline{\mathcal{B}}_{(1,6)(2,5)}$ so that the intersection is reducible and $\dim \mathcal{B}_{(1,3)(4,6)} = 2 \cdot 4 - 2 = 6$, $\dim \mathcal{B}_{(1,6)(2,5)} = 2 \cdot 4 - 4 = 4$.


5.2. As we explained in the introduction the meanders consisting of link patterns of orbital varieties that is M_{σ_S, σ_T} , where $S, T \in \mathbf{Tab}_{(n-k, k)^*}$ are the most simple on one hand and the most important on the other hand. To simplify the notation we put $P_T := P_{\sigma_T}$ and $M_{S, T} := M_{\sigma_S, \sigma_T}$.

By the very nature of the ordering of $\mathbf{B}$ orbits if $\mathcal{V}_T \cap \mathcal{V}_S$ is of higher (than 1) codimension and $\mathcal{B}_\sigma$ is a component of the intersection there exist $\mathcal{B}_\tau \subset \mathcal{V}_T$, $\mathcal{B}_\tau \not\subset \mathcal{V}_S$ and $\mathcal{B}_{\tau'} \subset \mathcal{V}_S$, $\mathcal{B}_{\tau'} \not\subset \mathcal{V}_T$ such that $\dim \mathcal{B}_\tau = \dim \mathcal{B}_{\tau'} = \dim \mathcal{B}_\sigma + 1$ and $\mathcal{B}_\sigma$ is a component of their intersection. Thus, the description of the intersections of higher codimension involves the description of intersections of codimension 1 for general $\mathbf{B}$ orbits. Just to illustrate this point let us show that the example of 5.1 provides

us with the intersection of codimension 2 of two orbital varieties which is not only reducible but also is not of pure dimension. This is the example from [13, 5.7]. Consider

$$T = \begin{array}{cc} 1 & 3 \\ 2 & 6 \\ 4 & \\ 5 & \end{array} \quad \text{and} \quad S = \begin{array}{cc} 1 & 2 \\ 3 & 5 \\ 4 & \\ 6 & \end{array}$$

so that $\sigma_T = (2, 3)(5, 6)$ and $\sigma_S = (1, 2)(4, 5)$. One has $\dim \mathcal{B}_{(2,3)(5,6)} = \dim \mathcal{B}_{(1,2)(4,5)} = 8$. The corresponding meander is



Comparing with the meander in 5.1 we see that $\mathcal{B}_{(1,3)(4,5)} \subset \overline{\mathcal{B}}_{(1,2)(4,5)}$ of codimension 1 and $\mathcal{B}_{(1,3)(4,5)} \not\subset \overline{\mathcal{B}}_{(2,3)(5,6)}$, also $\mathcal{B}_{(2,3)(4,6)} \subset \overline{\mathcal{B}}_{(2,3)(5,6)}$ of codimension 1 and $\mathcal{B}_{(2,3)(4,6)} \not\subset \overline{\mathcal{B}}_{(1,2)(4,5)}$. Moreover computing R_{σ_T, σ_S} we get

$$R_{\sigma_T, \sigma_S} = \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} = R_{(1,3)(4,5), (2,3)(4,6)}$$

Thus, exactly as in 5.1 we get

$$\mathcal{V}_T \cap \mathcal{V}_S = (\overline{\mathcal{B}}_{(1,3)(4,5)} \cap \overline{\mathcal{B}}_{(2,3)(4,6)}) \cap \mathcal{O}_{(4,2)^*} = (\overline{\mathcal{B}}_{(1,3)(4,6)} \cup \overline{\mathcal{B}}_{(1,6)(2,5)}) \cap \mathcal{O}_{(4,2)^*}$$

and $\dim \mathcal{B}_{(1,3)(4,6)} = 6$ and $\dim \mathcal{B}_{(1,6)(2,5)} = 4$ so that $\mathcal{V}_T \cap \mathcal{V}_S$ contains one component of codimension 2 and another component of codimension 4.

5.3. To formulate the statement on the intersections of codimension 1 we need more notation.

Arcs of $M_{S,T}$ never intersect one another. A connected subset of arcs in $M_{S,T}$ can be either open or closed. If it is open we call it an **interval**. If it is closed we call it a **loop**. The number of arcs in a path $\mathcal{P}$ is called the length of $\mathcal{P}$, we denote it by $L(\mathcal{P})$. Note that if $\mathcal{P}$ is a loop $L(\mathcal{P})$ is even. An interval can be either of even length, then we call it even, or of odd length, correspondingly we call it odd. We call a meander **even** if it consists only of loops and even intervals. Otherwise we call it **odd**. For example, the meander in 5.2 consists of two even intervals (each one of length 2) so it is an even meander.


Now we can formulate our result.

Theorem. *For $S, T \in \mathbf{Tab}_{(n-k, k)^*}$ one has $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) = 1$ if and only if $M_{S, T}$ is an even meander with $k - 1$ loops. In this case $\mathcal{F}_T \cap \mathcal{F}_S$ is irreducible.*

Proof. Note that $\text{codim}_{\mathcal{V}_T}(\mathcal{V}_T \cap \mathcal{V}_S) = 1$ iff $\mathcal{D}(\sigma_T) \cap \mathcal{D}(\sigma_S) \neq \emptyset$.


Since for $\sigma' \in D(\sigma_T)$ one has $\dim \mathcal{B}_{\sigma'} = k(n - k) - 1$ we get that either there is one fixed point under one arc of $P_{\sigma'}$ or exactly two arcs of $P_{\sigma'}$ intersect.

Let $P_{\sigma'}$ be a link pattern with one fixed point under one arc. Let m be this fixed point and (i, j) be the corresponding arc. There are exactly two ways to get a bigger link pattern (that is without fixed points under an arc and without crossings) and by maximality of dimension each one of them is a link pattern of some orbital variety:




where dots on $[i + 1, m - 1]$ are all end points of arcs belonging to $\pi_{i, m}(P_{\sigma'})$ and dots on $[m + 1, j - 1]$ are end points of arcs belonging to $\pi_{m, j}(P_{\sigma'})$.

Note that $M_{S, T}$ in that case consists of $k - 1$ loops $(i_s, j_s) \neq (i, m), (m, j)$ (all the loops are of length 2) and a unique even interval (of length 2) connecting i and j via m :




Now let $P_{\sigma'}$ have two intersecting arcs. Let $(i, j), (i', j')$ where $i < i' < j < j'$ be these arcs. There are exactly two ways to get a bigger link pattern (that is without crossings and without fixed points under an arc) and by maximality of dimension each one of them is a link pattern of some orbital variety:



where dots on $[i+1, i'-1]$ are all end points of arcs belonging to $\pi_{i,i'}(P_{\sigma'})$, dots on $[i'+1, j-1]$ are all end points of arcs belonging to $\pi_{i',j}(P_{\sigma'})$ and dots on $[j+1, j'-1]$ are all end points of arcs belonging to $\pi_{j,j'}(P_{\sigma'})$.

Note that $M_{S,T}$ consists of $k-1$ loops: $k-2$ loops of type (i_s, j_s) (of length 2) where $\{i_s, j_s\} \cap \{i, i', j, j'\} = \emptyset$ and a loop (of length 4) involving $\{i, i', j, j'\}$:



Thus, in both cases we get an even meander with $k-1$ loops. Let $M_{S,T}$ be a meander with $2k$ arcs. If it is even and contains $k-1$ loops then it must contain $k-2$ loops of length 2 and either one loop of length 4 or one more loop of length 2 and one even interval of length 2, so that the cases above are all possible cases for even meanders with $k-1$ loops.

Also, one can see from the pictures above that for $\sigma \in \mathcal{D}(\sigma_T)$ there exists exactly one S such that $\mathcal{D}_{\sigma_T} \cap \mathcal{D}_{\sigma_S} \supset \{\sigma\}$ and in this case $\mathcal{D}_{\sigma_T} \cap \mathcal{D}_{\sigma_S} = \{\sigma\}$ so that the intersection of codimension 1 is irreducible. $\square$

5.4. Let us compare the results above to the results of Wesbury [18] which were applied by Fung [2] to two-row case. In this subsection we formulate the results which coincide with our results on two-column case.

For $T \in \mathbf{Tab}_{(n-k,k)}^*$ let $I(T) := I(P_{\sigma_T}) := I(\sigma_T) := \{i : (i, i+1) \in \sigma_T\} = \{i : i \in \langle T_1 \rangle, i+1 \in \langle T_2 \rangle\}$. For example, take

$$T = \begin{array}{cc} 1 & 4 \\ 2 & 5 \\ 3 & 7 \\ 6 & 8 \\ 9 & \end{array}$$

Then $\sigma_T = (3, 4)(2, 5)(6, 7)(1, 8)$ and $I(T) = \{3, 6\}$.

Given $i \in \langle T_1 \rangle$, $j \in \langle T_2 \rangle$ put $T_{i \rightleftharpoons j}$ to be a tableau obtained from T by interchanging columns of i and j if an array obtained in such a way is a tableau. Otherwise $T_{i \rightleftharpoons j} = \emptyset$. Note that if $j < i$ then $T_{i \rightleftharpoons j} \neq \emptyset$ always. If $j > i$ then $T_{i \rightleftharpoons j} \neq \emptyset$ iff $\text{sh}(\pi_{1,i}(T)) = (i - k', k')^*$ where $i - k' > k' + 1$ and $\text{sh}(\pi_{1,j}(T)) = (j - k'', k'')^*$ where $j - k'' > k''$.

If $i \notin I(T)$ put


$$u_i(T) := \begin{cases} T_{\sigma_T(i) \rightleftharpoons i} & \text{if } i, i+1 \in \langle T_2 \rangle \\ T_{i+1 \rightleftharpoons i} & \text{if } i \in \langle T_2 \rangle \text{ and } i+1 \in \langle T_1 \rangle \\ T_{i+1 \rightleftharpoons \sigma_T(i+1)} & \text{if } i, i+1 \in \langle T_1 \rangle \text{ and } \sigma_T(i+1) \neq i+1 \\ \emptyset & \text{if } i, i+1 \in \langle T_1 \rangle \text{ and } \sigma_T(i+1) = i+1 \end{cases}$$

Note that in the last case $\sigma_T(i) = i$ as well. And this is the only case when $u_i(T) = \emptyset$.


The picture is much more clear on the level of link patterns. Let T be such that $i \notin I(P_T)$. then $u_i(P_T) \neq \emptyset$ iff $\{i, i+1\} \cap \langle P_T \rangle \neq \emptyset$.

In this case


- If $\{i, i+1\} \subset \langle \sigma_T \rangle$ then $u_i(\sigma_T) := \sigma_{u_i(T)} = (\sigma_T)_{i \rightleftharpoons \sigma_T(i+1)}$ and $M_{T, u_i(T)}$ consists of $k - 2$ loops of length 2 and one loop of length 4 with the points $\{i, i+1, \sigma_T(i), \sigma_T(i+1)\}$. There are 3 possible situations shown below:



- If $\{i, i+1\} \not\subset \langle \sigma_T \rangle$ then
 - If $i \in \langle \sigma_T \rangle$ and $i+1 \notin \langle \sigma_T \rangle$ then $i \in \langle T_2 \rangle$, $i+1 \in \langle T_1 \rangle$ and $u_i(\sigma_T) := \sigma_{u_i(T)} = (\sigma_T)_{\sigma_T(i) \rightarrow i+1}$, so that $M_{T, u_i(T)}$ consists of $k-1$ loops of length 2 and one interval of length 2 consisting of arcs $(\sigma_T(i), i)$, $(i, i+1)$ as it is shown at the picture below:



- If $i+1 \in \langle \sigma_T \rangle$ and $i \notin \langle \sigma_T \rangle$ then $i, i+1 \in \langle T_1 \rangle$ and $u_i(\sigma_T) := \sigma_{u_i(T)} = (\sigma_T)_{\sigma_T(i+1) \rightarrow i}$ so that $M_{T, u_i(T)}$ consists of $k-1$ loops of length 2 and an interval of length 2 consisting of arcs $(i, i+1)$, $(i+1, \sigma_T(i+1))$ as it is shown at the picture below:



Note that in both cases we get by Theorem 5.3 that $\text{codim}_{\mathcal{V}_T}(\mathcal{V}_T \cap \mathcal{V}_{u_i(T)}) = 1$. Moreover, if $i \notin I(T)$ then $u_i(T)$ is the only tableau in $\{S \in \mathbf{Tab}_{(n-k,k)^*} : i \in I(S)\}$ such that $\text{codim}_{\mathcal{V}_T}(\mathcal{V}_T \cap \mathcal{V}_S) = 1$.

Further, by [18, 7] one can define inner product on Temperley-Lieb algebra of $\{P_T : T \in \mathbf{Tab}_{(n-k,k)}\}$ where link pattern P_T for $T \in \mathbf{Tab}_{(n-k,k)}$ is defined exactly in the same way as it is defined here (that is $P_T := P_{T^t}$). Then for two basis elements $P_T, P_S : T, S \in \mathbf{Tab}_{(n-k,k)}$ one has either $\langle P_T, P_S \rangle = \delta^s$ where $0 \leq s \leq k-1$ or $\langle P_T, P_S \rangle = 0$. The W graph of link patterns with vertices $\{P_T : T \in \mathbf{Tab}_{(n-k,k)}\}$ each labeled by the set $I(T)$ (where $i \in I(T)$ if i is in the first row of T and $i+1$ in the second row of T , that is $I(T) := I(T^t)$) and edges connecting P_T and P_S if $\langle P_T, P_S \rangle = \delta^{k-1}$ is a graph of Kazhdan-Lusztig type. Moreover, translating the result to our language we get $\langle P_{S^t}, P_{T^t} \rangle = \delta^{k-1}$ iff $M_{S,T}$ is an even meander with $k-1$ loops.

Comparing to our results we get that for $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ one has $\text{codim}_{\mathcal{F}_S}(\mathcal{F}_T \cap \mathcal{F}_S) = 1$ iff $\langle P_{S^t}, P_{T^t} \rangle = \delta^{k-1}$. Thus, in particular W graph of link patterns with vertices $\{P_T : T \in \mathbf{Tab}_{(n-k,k)^*}\}$ labeled by the set $I(T)$ and edges connecting P_T and P_S if $\text{codim}_{\mathcal{V}_S}(\mathcal{V}_T \cap \mathcal{V}_S) = 1$ coincides with the graph of corresponding

Templerley-Lieb algebra and respectively it is of Kazhdan-Lusztig type. As it was shown by F. Fung $\text{codim}_{\mathcal{F}_{S^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t}) = 1$ exactly in the same cases.

5.5. More generally, in [18] and [3] it was shown that for $P_S, P_T : S, T \in \mathbf{Tab}_{(n-k,k)}$

$$\langle P_S, P_T \rangle = \begin{cases} \delta^r & \text{if } M_{S,T} \text{ is an even meander with } r \text{ loops;} \\ 0 & \text{if } M_{S,T} \text{ is odd.} \end{cases}$$

They showed this theorem using the proposition that for $P_S, P_T : S, T \in \mathbf{Tab}_{(n-k,k)}$ such that $i \in I(P_S)$ and $i \notin I(P_T)$ one has

$$\langle P_S, P_T \rangle = \begin{cases} \delta^{-1} \langle P_S, P_{u_i(T)} \rangle & \text{if } u_i(P_T) \neq \emptyset; \\ 0 & \text{otherwise.} \end{cases}$$

F. Fung in [2] considered the intersections of $\mathcal{F}_T : T \in \mathbf{Tab}_{(n-k,k)}$. By [2, 7] for $T, S \in \mathbf{Tab}_{(n-k,k)}$ one has

$$\text{codim}_{\mathcal{F}_S}(\mathcal{F}_T \cap \mathcal{F}_S) = \begin{cases} k - r & \text{if } \langle P_S, P_T \rangle = \delta^r \\ \emptyset & \text{if } \langle P_S, P_T \rangle = 0 \end{cases}$$

He showed this using the same technique, namely showing that for $i \in I(P_S)$, $i \notin I(P_T)$ one has

$$\text{codim}_{\mathcal{F}_S}(\mathcal{F}_T \cap \mathcal{F}_S) = \begin{cases} \text{codim}_{\mathcal{F}_S}(\mathcal{F}_S \cap \mathcal{F}_{u_i(T)}) + 1 & \text{if } u_i(P_T) \neq \emptyset; \\ \emptyset & \text{otherwise.} \end{cases}$$

However the results on the intersections of higher codimensions in two-column case are very different from those of Westbury and Fung. First of all note that since $\mathcal{F}_T \cap \mathcal{F}_S \neq \emptyset$ for $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ it cannot occur that $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) = \text{codim}_{\mathcal{F}_{T^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t})$ for all $S, T \in \mathbf{Tab}_{(n-k,k)^*}$.

It is interesting that the deviations of the codimensions are directed in both sides. On one hand, if $M_{S,T}$ is odd then $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) < \text{codim}_{\mathcal{F}_{T^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t})$ (since $\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t} = \emptyset$). On the other hand, for even meanders with less than $k - 1$ loops it seems that $\text{codim}_{\mathcal{F}_T}(\mathcal{F}_T \cap \mathcal{F}_S) \geq \text{codim}_{\mathcal{F}_{T^t}}(\mathcal{F}_{T^t} \cap \mathcal{F}_{S^t})$. We do not prove this here, we only show by the example that the technique of passing from $M_{T,S}$ to $M_{u_i(T),S}$ explained above does not work in two-column case. Namely we give an example of $S, T \in \mathbf{Tab}_{(n-k,k)^*}$ such that $i \in I(S)$ and $i \notin I(T)$ and $u_i(T) \neq \emptyset$ and $\text{codim}_{\mathcal{F}_S}(\mathcal{F}_T \cap \mathcal{F}_S) > \text{codim}_{\mathcal{F}_S}(\mathcal{F}_{u_i(T)} \cap \mathcal{F}_S) + 1$:

Let us consider $S, T \in \mathbf{Tab}_6^2$ shown below. Note that $2 \in I(S)$ and $2 \notin I(T)$ so that

$$S = \begin{array}{cc} 1 & 3 \\ 2 & 4 \\ 5 & \\ 6 & \end{array}, \quad T = \begin{array}{cc} 1 & 5 \\ 2 & 6 \\ 3 & \\ 4 & \end{array}, \quad u_2(T) = \begin{array}{cc} 1 & 3 \\ 2 & 5 \\ 4 & \\ 6 & \end{array}$$

The corresponding meanders are:



By Theorem 5.3 $\text{codim}_{\mathcal{F}_S} \mathcal{F}_S \cap \mathcal{F}_{u_2(T)} = 1$. On the other hand, a straightforward computation shows that $\mathcal{V}_S \cap \mathcal{V}_T$ is irreducible and its only component is $\mathcal{B}_{(1,6)(2,5)}$ with $\dim \mathcal{B}_{(1,6)(2,5)} = 8 - 4 = 4$ so that $\text{codim}_{\mathcal{F}_S} \mathcal{F}_S \cap \mathcal{F}_T = \text{codim}_{\mathcal{F}_S} \mathcal{F}_S \cap \mathcal{F}_{u_2(T)} + 3$.

One can define $I(\sigma)$ and respectively $u_i(\sigma)$ for any σ not only for σ_T but in general case $\dim \mathcal{B}_{u_i(\sigma)} \geq \dim \mathcal{B}_\sigma$ and the difference between these two dimensions can be quite big. And this is the reason of the jumps of codimension of the intersections. But the details of this general theory are very involved on one hand and $\mathcal{F}_T \cap \mathcal{F}_S$ of higher codimensions are not that interesting from the geometric point of view on the other hand, so we stop here.

5.6. Let us note at the end that there exists a very easy sufficient condition for $\mathcal{F}_T \cap \mathcal{F}_S$ to be reducible. Generalizing this condition we can obtain the necessary and sufficient condition for the intersection to be irreducible, but again, the formulation of this condition is too involved.

Consider $M_{S,T}$ and put $[i_1, j_1], [i_2, j_2], \dots$ to be the segments such that $(R_{S,T})_{i_s, j_s} = 1$ and $(R_{S,T})_{p,q} = 0$ for any $[p, q] \subsetneq [i_s, j_s]$. We order these segments in increasing order (that is $i_1 < i_2 < \dots$) and call them 1-segments of $M_{S,T}$.

Proposition. *Consider $S, T \in \mathbf{Tab}_{(n-k,k)^*}$. Let $[i_1, j_1], \dots, [i_t, j_t]$ be 1-segments of $M_{S,T}$. If there exists $s : 1 \leq s < t$ such that either $i_{s+1} = j_s$ or $i_{s+1} < j_s$ and $(R_{S,T})_{i_s, j_{s+1}} = 1$, then $\mathcal{F}_S \cap \mathcal{F}_T$ is reducible.*

Proof. First of all note that for a 1-segment $[i_s, j_s]$ there exists a component $\mathcal{B}_\sigma$ of $\overline{\mathcal{B}}_{\sigma_S} \cap \overline{\mathcal{B}}_{\sigma_T}$ such that $(i_s, j_s) \in \sigma$. Indeed, by the definition of $[i_s, j_s]$ one has $(i_s, j_s) \prec \sigma_S, \sigma_T$. Thus, by proposition 3.10 there exists a component involution σ of $\overline{\mathcal{B}}_{\sigma_S} \cap \overline{\mathcal{B}}_{\sigma_T}$ such that $\sigma \succ (i_s, j_s)$. In particular, this provides $(R_\sigma)_{i_s, j_s} \geq 1$. On the other hand, for any $[p, q] \subsetneq [i_s, j_s]$ one has $(R_\sigma)_{p,q} \leq (R_{S,T})_{p,q} = 0$ and $(R_\sigma)_{i_s, j_s} \leq (R_{S,T})_{i_s, j_s} = 1$. Thus, $(R_\sigma)_{i_s, j_s} = 1$ and $(i_s, j_s) \in \sigma$.

If $i_{s+1} = j_s$ then there exist component involutions σ, σ' of $\mathcal{B}_T \cap \mathcal{B}_S$ such that $(i_s, j_s) \in \sigma$ and $(j_s, j_{s+1}) \in \sigma'$. Obviously, $\mathcal{B}_\sigma, \mathcal{B}_{\sigma'}$ are two different components of the intersection.

If $i_{s+1} < j_s$ then there exist component involutions σ, σ' such that $(i_s, j_s) \in \sigma$ and $(j_s, j_{s+1}) \in \sigma'$. If $\sigma = \sigma'$ we get $(i_s, j_s), (i_{s+1}, j_{s+1}) \in \sigma$ so that $(R_\sigma)_{i_s, j_{s+1}} = 2$ which contradicts the condition $(R_{S,T})_{i_s, j_{s+1}} = 1$. Thus, again, $\mathcal{B}_\sigma, \mathcal{B}_{\sigma'}$ are two different components of the intersection. $\square$

INDEX OF NOTATION

- 1.1 $\mathcal{O}_u, \mathfrak{N}, \mathbf{B}, \mathcal{B}_u, \mathcal{X}$
- 1.2 $\lambda, \lambda^*, J(u), \mathcal{O}_\lambda, D_\lambda$
- 1.3 $\mathbf{S}_n^2, N_\sigma, \mathcal{B}_\sigma, P_\sigma$
- 1.4 $\ell(P_\sigma), c(P_\sigma), f_p(P_\sigma), f(P_\sigma), \mathbf{S}_n^2(k)$
- 1.5 $[i, j], R_\sigma, \succeq$
- 1.6 $\mathcal{C}(\sigma), \mathcal{N}(\sigma), \mathcal{D}(\sigma)$
- 1.9 $R_{\sigma, \sigma'}$
- 2.1 $\mathfrak{B}_{(n-k, k)}, (i, j), ((i, j)), q_{(i, j)}(\sigma)$
- 2.2 $\mathcal{V}_T, \mathbf{Tab}_{(n-k, k)^*}, T_1, T_2, \langle T_i \rangle, \sigma_T, \mathcal{B}_T$
- 2.3 $\pi_{i, j}, R_u, R_\sigma, \mathbf{R}_n^2$
- 3.1 $\mathbf{over}_m(P_\sigma), \mathbf{over}_{(i, j)}(P_\sigma), l_{(i, j)}(P_\sigma), r_{(i, j)}(P_\sigma), c_{(i, j)}^l(P_\sigma),$
 $c_{(i, j)}^r(P_\sigma), \mathbf{under}_{(i, j)}(P_\sigma), f'_{(i, j)}(P_\sigma)$
- 3.2 $\pi_{i, j}(P_\sigma)$
- 3.3 $E(\sigma), \sigma_{(i, j)}^-$
- 3.4 $\langle P_\sigma \rangle, \langle \sigma \rangle, \sigma_{i \rightarrow f}, \sigma_{\curvearrowright (i, j)}, \sigma_{(i, j) \curvearrowright}$
- 3.5 $\sigma_{i \rightleftharpoons j}, L_{(i, j)}(\sigma), S_{(i, j) \rightleftharpoons}$
- 3.6 $ov_{(i, j)}(\sigma), S_{(i, j) \downarrow \uparrow}$
- 3.8 $E_{\max}(\sigma)$

REFERENCES

- [1] P. Di Francesco, P. Zinn-Justin, *From Orbital Varieties to Alternating Sign Matrices*, math-ph/0512047.
- [2] F.Y.C.Fung *On the topology of components of some Springer fibers and their relation to Kazhdan-Lusztig theory*, Advances in Math. 178 (2003), pp. 244-276.
- [3] J.J. Graham, G.I. Lehrer, *Cellular algebras*, Invent. Math. **123** (1996), pp. 1-34.
- [4] W. Hesselink, *Singularities in the nilpotent scheme of a classical group*, Trans. Am. Math. Soc. **222** (1976), pp. 1-32.
- [5] A. Joseph, *On the variety of a highest weight module*, J. of Algebra 88 (1984), 238-278.
- [6] A. Knutson, P. Zinn-Justin, *A scheme related to the Brauer loop model*, math.AG/0503224.
- [7] A.Melnikov, *Irreducibility of the associated varieties of simple highest weight modules in $\mathfrak{sl}(n)$* , C.R.A.S. I, 316 (1993), 53-57.
- [8] A.Melnikov, *Orbital varieties in $\mathfrak{sl}_n$ and the Smith conjecture*, J. of Algebra, **200** (1998), pp. 1-31.
- [9] A. Melnikov, *B-orbits in solutions to the equation $X^2 = 2$ in triangular matrices*, J. of Algebra, **223** (2000), pp. 101-108.
- [10] A.Melnikov,, *The combinatorics of orbital variety closures of nilpotent order 2 in $\mathfrak{sl}_n$* , Electronic Journal of Combinatorics, vol. 12(1), 2005, R21.
- [11] A. Melnikov, *Description of B-orbit closures of order 2 in upper-triangular matrices*, Transformation Groups, Vol. 11 No. 2, 2006, pp. 217-247.
- [12] A. Melnikov, N.G.J. Pagnon, *On the intersections of orbital varieties and components of Springer fiber*, J. Algebra 298 (2006), 1-14.

- [13] A. Melnikov, J.G.N. Pagnon, *Intersections of components of a Springer fiber for the hook and two column cases*, math.RT/0607673.
- [14] A. Melnikov, J.G.N. Pagnon, *Intersections of components of a Springer fiber of codimension one for the two column case*, math.RT/0701178.
- [15] N.Spaltenstein, *The fixed point set of a unipotent transformation on the flag manifold*, Proc. Konin. Nederl. Akad. **79** (1976), pp. 452-456.
- [16] N. Spaltenstein, *Classes unipotentes de sous-groupes de Borel*, LN in Math. 964 Springer-Verlag, 1982.
- [17] R. Steinberg, *On the desingularization of the unipotent variety*, Invent. Math. 36 (1976), 209-224.
- [18] B.W. Westbury, *The representation theory of the Temperley-Lieb algebras*, Math. Zeitschrift, 219 (1995), pp. 539-565.
- [19] J.-B. Zuber, *On the counting of fully packed loop configurations: some new conjectures*, The Electronic Journal of Combinatorics, 11(2004), R13

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA 31905, ISRAEL
E-mail address: melnikov@math.haifa.ac.il